



This is a digital copy of a book that was preserved for generations on library shelves before it was carefully scanned by Google as part of a project to make the world's books discoverable online.

It has survived long enough for the copyright to expire and the book to enter the public domain. A public domain book is one that was never subject to copyright or whose legal copyright term has expired. Whether a book is in the public domain may vary country to country. Public domain books are our gateways to the past, representing a wealth of history, culture and knowledge that's often difficult to discover.

Marks, notations and other marginalia present in the original volume will appear in this file - a reminder of this book's long journey from the publisher to a library and finally to you.

Usage guidelines

Google is proud to partner with libraries to digitize public domain materials and make them widely accessible. Public domain books belong to the public and we are merely their custodians. Nevertheless, this work is expensive, so in order to keep providing this resource, we have taken steps to prevent abuse by commercial parties, including placing technical restrictions on automated querying.


We also ask that you:

- + *Make non-commercial use of the files* We designed Google Book Search for use by individuals, and we request that you use these files for personal, non-commercial purposes.
- + *Refrain from automated querying* Do not send automated queries of any sort to Google's system: If you are conducting research on machine translation, optical character recognition or other areas where access to a large amount of text is helpful, please contact us. We encourage the use of public domain materials for these purposes and may be able to help.
- + *Maintain attribution* The Google "watermark" you see on each file is essential for informing people about this project and helping them find additional materials through Google Book Search. Please do not remove it.
- + *Keep it legal* Whatever your use, remember that you are responsible for ensuring that what you are doing is legal. Do not assume that just because we believe a book is in the public domain for users in the United States, that the work is also in the public domain for users in other countries. Whether a book is still in copyright varies from country to country, and we can't offer guidance on whether any specific use of any specific book is allowed. Please do not assume that a book's appearance in Google Book Search means it can be used in any manner anywhere in the world. Copyright infringement liability can be quite severe.

About Google Book Search


Google's mission is to organize the world's information and to make it universally accessible and useful. Google Book Search helps readers discover the world's books while helping authors and publishers reach new audiences. You can search through the full text of this book on the web at <http://books.google.com/>

Ced
W56






Ced
W56











CATALOGUS

PLANTARUM MEDICINALIUM.

Ced
W56

CATALOGUS RATIONALIS

PLANTARUM MEDICINALIUM,

IN HORTO

SOCIETATIS PHARMACEUTICÆ LONDINENSIS,

APUD VICUM CHELSEA, CULTARUM.

CURANTE

JACOBO L. WHEELER,
SOCIETATIS BOTANICES PROFESSORE,
SOC. LINN. SOCIO.
ET IN CHEMIA ET MATERIA MEDICA PRELECTORE.

LONDINI,
EX TYPOGRAPHIA GULIELMI TYLER,
BRIDGEWATER SQUARE.

1830.

Ced

W56

D. JOSEPHO HURLOCK,

MAGISTRO,

D. GULIELMO RIO MACDONALD,

D. JOHANNI HUNTER,

CUSTODIBUS,

CÆTERISQUE CURATORIBUS,

SOCIETATIS PHARMACEUTICÆ LONDINENSIS,

HOC OPUSCULUM,

IN USUM JUVENTUTIS STUDIOSE,

EORUM JUSSU SUSCEPTUM,

SUMMI HONORIS CAUSA,

DICAT, DEDICATQUE

EDITOR.

CATALOGUS, &c.

CLASSIS I.

ORDO I.

MONANDRIA. MONOGYNIA.

* *Scitamineæ fructu infero uni—vel triloculari.*

CURCUMA. Roscoe, Actis Societatis Linneanæ, tom. viii. 354.

ORDO NATURALIS SCITAMINA. Linn.

CLASSIS IV. PLANTÆ MONOCOTYLEDONES.
STAMINA EPIGYNA. ORDO II. CANNÆ. Jussieu
Genera Plantarum.

CHAR. GEN. Anthera duplex, bicalcarata. Filamentum petaliforme, trilobatum, medio antheriferum.

1. CURCUMA ZEDOARIA.

CHAR. SPEC. Foliis ovatis acuminatis, bracteis emarginatis.

Amomum Zedoaria. Willdenow Species Plantarum, tom. i. 7.

Amomum Zedoaria. Persoon Synopsis, tom. i. 2.

ANGLICE, Zedoary.

LOCUS. Zeylona, Malabar. Perennis.

VIS. Subacris aromatica.

2. CURCUMA LONGA. Willd. Sp. Pl. i. 14. Persoon Syn. i. 4. Roscoe, Act. Soc. Linn. tom. viii. 355.

CHAR. SPEC. Foliis ovato-lanceolatis, bracteis spatulatis.

ANGLICE, Turmeric.

LOCUS. Amboina, Malabaria, Zeylona. Perennis.

QUALITAS. Amara, luteo tinctoria.

ALPINIA. Willd. Spec. Plant. i. 11. Persoon Synopsis Plantarum, tom. i. 3.

ORDO NAT. SCITAMINA. Linn.

CLASSIS IV. ORDO II. CANNÆ. Juss.

CHAR. GEN. Calyx tridentatus æqualis tubulosus. Corolla tripartita æqualis. Nectarium bilabiatum labio inferiore patente.

3. ALPINIA CARDAMOMUM.

CHAR. SPEC. Foliis lanceolatis scapis basilaribus flexuosis, labello trilobo basi calcarato. Roxburgh Flora Indica, tom. i. 68.

Elettaria Cardamomum. Maton, Actis Societat. Linn. x. part. ii. p. 254.

Amomum Cardamomum. Willd. Sp. Plant. tom. i. 8. Persoon Synopsis, tom. i. 2.

ANGLICE, Lesser Cardamom.

LOCUS. Malabaria, Zeylona. Perennis.

VIS. Calefaciens, stimulans.

COMP. Tinct. Cardam. *Phar. Lond. Edin. Eblan.* Tinct. Card. Comp. *Phar. Lond. Eblan.* et alia plura.

ZINGIBER. Act. Soc. Linn. viii. 347.

ORDO NAT. SCITAMINA. Linn.

CANNÆ. Juss.

CHAR. GEN. Anthera duplex. Filamentum extra antheram elongatum; apice subulatum, sulcatum. Stylus in sulco antheræ receptus.

4. ZINGIBER OFFICINALE.

CHAR SPEC. Bracteis ovato-lanceolatis, laciniis corollæ revolutis, nectario trilobato. Amomum Zingiber. Willd. Sp. Pl. i. 6. Persoon Syn. i. 2.

ANGLICE, Ginger.

LOCUS. India Orientalis, Jamaica, et Indiæ occidentalis ceteræ insulæ. Perennis.

VIS. Acris aromatica.

USUS. Dyspepsiâ, colicâ. Purgantia corrigens.

DOSIS. Gr. x. ad ʒi.

COMPOSITA. Syr. Zingiberis. *Phar. Lond. Edin. Eblan.* Tinct. Zingiberis. *Ph. Lond. Eblan.* et alia plura.

MARANTA. Roscoe, Actis Soc. Linn. tom. viii. 339.

SCITAMINA. Linn.

CANNÆ. Juss.

CHAR. GEN. Anthera simplex, filamentum adnatum. Stylus petaliformis. Stigma subtrigonum. Semen unicum.

5. MARANTA ARUNDINACEA.

CHAR. SPEC. Culmo ramoso herbaceo, foliis ovato-lanceolatis subtus pilosiusculis. Willd. Sp. Pl. i. 13. Persoon Syn. i. 3.

ANGLICE, Arrow-root.

LOCUS. America Meridionalis. Perennis.

Radix amylo abundat, quod aquâ bulliente facillime liquifacit, victum tenuem ægrotis præstat.

CLASSIS II.

ORDO I.

DIANDRIA. MONOGYNIA.

* *Flores inferi, Monopetali, Regulares.*

OLEA. Willd. Spec. Plant. i. 44. Persoon Syn. i. 8.

ORDO NAT. SEPIARLÆ. Linn.

**CLASSIS VIII. PLANTÆ DICOTYLEDONES
MONOPETALÆ. COROLLA HYPOGYNA. ORDO
IV. JASMINEÆ.**

II. Fructus baccatus. Juss.

CHAR. GEN. Corolla quadrifida laciniis subovatis.
Drupa disperma : uno abortante.

6. OLEA EUROPÆA.

CHAR. SPEC. Foliis lanceolatis integerrimis, racemis axillaribus coarctatis.

ANGLICE, European Olive.

LOCUS. Hispania, Gallia, Italia. Perennis.

Oleum e drupis exprimitur, inodorum, et insipidum : cujus pondus specificum est. 960. Congelat circa gradum 38vum. et in elementa resolvitur ad gradum 600. Sæpenumero oleo papaverum fraudulentè commiscetur.

In emulsionibus, emplastris, linimentis, conficiendis adhibetur.

**** Flores inferi, monopetali, irregulares. Fructus capsularis.**

VERONICA. Willd. Sp. Pl. i. 54. Persoon Syn. i. 10.

**ORDO NAT. PERSONATÆ. Linn. CLASSIS
VIII. ORDO II. PEDICULARES.**

I. Stamina non didynama, duo aut plura. Juss.

CHAR. GEN. Corolla quadrifida, limbo planiusculo, lacinia inferiore angustiore. Capsula bilocularis.

**** Corymboso-racemosæ.**

7. — 8. VERONICA BECCABUNGA.

CHAR. SPEC. Racemis lateralibus, foliis ovatis, planis, caule repente.

ANGLICE, Brooklime.

LOCUS. Europæ scaturigines. Perennis.

VIS. Antiscorbutica olim existimata, nunc inusitata.

GALIPEA. Willd. Sp. Pl. i. 41. Persoon Syn. i. 8.

**ORDO NAT. PLANTÆ INCERTÆ SEDIS. MO-
NOPETALÆ GERMINE SUPERO. Juss.**

CHAR. GEN. Calyx tetragonus vel pentagonus quadri- vel quinquedentatus. Corolla hypocrateriformis profundè 4 vel 5 partita. Stamina quatuor : duo sterilia.

8. ~~7.~~ **GALIPEA CUSPARIA.** Plantæ Aequinoctiales, Vol. ii. p. 59, t. 57.

CHAR. SPEC. Foliis ternatis racemis pedunculatis, subterminalibus calyce quinquedentato staminibus sterilibus tribus.

ANGLICE, Cusparia.

LOCUS. Ad fluviorum ripas Guianæ. Perennis.

VIS. Stimulans, tonica in febribus intermittentibus, diarrhœâ, dyspepsiâ.

COMP. Inf. cuspariæ Lond. Tinctura. Edin. et Eblan.

GRATIOLA. Willd. Spec. Plant. i. 102. Persoon Syn. i. 14.

ORDO NAT. PERSONATÆ. Linn.

CLASSIS VIII. ORDO VII. SCROPHULARIÆ.

III. Genera scrophulariis affinia, oppositifolia. Juss.

CHAR. GEN. Calyx heptaphyllus, exterior patulus, corolla resupinata. Filamenta duo sterilia. Capsula bilocularis.

9. GRATIOLA OFFICINALIS.

CHAR. SPEC. Foliis lanceolatis, serratis, floribus pedunculatis.

ANGLICE, Hedge hyssop.

LOCUS. Europa Australior. Perennis.

VIS. Amara, cathartica, emetica, diuretica : in Germania usitata.

*** Flores inferi, monopetali, irregulares. Fructus gymnospermi.

ROSMARINUS. Willd. Spec. Plant. i. 126. Persoon Syn. i. 24.

ORDO NAT. VERTICILLATÆ. Linn.

CLASSIS VIII. ORDO VI. LABIATÆ. I. Stamina
duo fertilia, duo abortiva. Juss.

CHAR. GEN. Corolla ringens, labio superiore bipartito.
Filamenta longa, curva simplicia, cum dente.

10. ROSMARINUS OFFICINALIS.

ANGLICE, Rosemary.

LOCUS. Gallia Narbonensis, Hispania. Perennis.

VIS. Aromatica, pungens.

Emmenagoga olim existimata.

SALVIA. Willd. Spec. Plant. i. 127. Persoon Syn.
i. 24.

ORDO NAT. VERTICILLATÆ. Linn.

CLASSIS VIII. ORDO VI. LABIATÆ. Juss.

CHAR. GEN. Corolla ringens. Filamenta transversè
pedicello affixa.

11. SALVIA OFFICINALIS.

CHAR. SPEC. Foliis lanceolato-ovatis crenulatis, ver-
ticillis paucifloris, calycibus mucronatis.

ANGLICE, Sage.

LOCUS. Italia, Narbona. Perennis.

VIS. Amara Aromatica.

CLASSIS II.

ORDO III.

DIANDRIA. TRIGYNIA.

PIPER. Willd. Spec. Plant. i. 159. Persoon Syn.
i. 31.

ORDO NAT. PIPERITÆ. Linn.

**CLASSIS XV. PLANTÆ DICOTYLEDONES
APETALÆ. STAMINA IDIOGYNA, SEU A PIS-
TILLO SEGREGATA.**

ORDO III. URTICÆ. III. Genera urticis affinia. Juss.

CHAR. GEN. Spadix amentiformis. Calyx nullus.
Corolla nulla. Stylus brevissimus. Stigmata tria, mani-
festa. Bacca monosperma.

12. PIPER NIGRUM.

CHAR. SPEC. Foliis ovatis subseptemnerviis glabris,
petiolis simplicibus.

ANGLICE, Black pepper.

LOCUS. India Orientalis, Malacca, Java, Sumatra.
Perennis.

VIS. Acris.

DOSIS. Gr. x. ad ʒi.

COMP. Ung. Piperis Nig. *Pharm. Eblan.* Emp. Meloës
Vesicator, Comp. Edin. Conf. Piper. Nigri. Lond.

Nuperrime compertum est a D. Pelletier, Piper nigrum ma-
teriam propriam, valde acrem, et Piperin nuncupatam, sed
alkalibus minime affinem, continere.

13. 14. PIPER CUBEBA. Willd. Sp. Pl. i. 159. Per-
soon Syn. i. 31.

CHAR. SPEC. Foliis oblique ovatis, oblongis, venosis,
acutis, spica solitaria pedunculata oppositifolia, fructibus
pedicellatis.

ANGLICE, Cubebs.

LOCUS. Javæ inque Guineæ sylvis.

VIS. Aromatica, amara.

DOSIS. ʒss ad ʒij quater die.

USUS. Gonorrhœa. In Societatis Med-Chirurgicæ
Actis, tom. xii. D. Broughton sequentem edidit exposi-
tionem, hujus medicamenti potestatem optime illustrantem,
hominum quinquaginta sub Gonorrhœâ laborantium, qui hoc
remedium tentaverunt,
Intra septem dies sani evaserunt decem.
Intra quatuordecim dies, septendecim.

Intra triginta dies, unus.

Intra quinquaginta quinque dies, unus.

At nauseam, cephalalgiam, vertiginem aliquando provocat.

14. ~~13.~~ PIPER LONGUM. Willd. Sp. Plant. i. 161. Persoon Syn. i. 31.

CHAR. SPEC. Foliis cordatis petiolatis sessilibusque.

ANGLICE, Long Pepper.

LOCUS. Malabaria, Bengalia. Perennis.

CLASSIS III.

ORDO I.

TRIANDRIA. MONOGYNIA.

* *Flores Superi.*

ORDO NAT. AGGREGATÆ. Linn.

CLASSIS XI. DICOTYLEDONES MONOPETALÆ.

COROLLA EPIGYNA. ANTHERÆ DISTINCTÆ.

ORDO I. DIPSACÆ. II. Flores distincti. Juss.

VALERIANA. Willd. Spec. Plant. i. 175. Persoon Syn. i. 37.

CHAR. GEN. Calyx nullus, aut margo minimus. Corolla monopetala, subinfundibuliformis, quinquefida, basi hinc gibba, supera. Semen unicum.

15. VALERIANA OFFICINALIS.

CHAR. SPEC. Foliis omnibus pinnatis, foliolis dentatis lanceolatis, caule sulcato.

ANGLICE, Valerian.

LOCUS. Europæ prata paludosa. Perennis.

VIS. Amara, antispasmodica?

USUS. Hysteriâ, Epilepsiâ.

DOSIS. Radicis pulveris ʒi ad ʒi quater die.

COMP. Ext. Valerianæ, *Phar. Eblan.* Inf. Valerian, *Phar. Eblan.* Tr. Valerian, *Phar. Lond. et Eblan.* Tr. Val. Ammoniata. *Lond. Edin. et Eblan.*

CROCUS. Willd. Sp. Pl. i. 194. Persoon Syn. i. 41.

ORDO NAT. DENUDATÆ. Linn.

CLASSIS III. PLANTÆ MONOCOTYLEDONES.
STAMINA PERIGYNA.

ORDO VIII. IRIDES. II. Stamina filamentis distinctis.
Juss.

CHAR.GEN. Spatha radicalis. Corolla infundibuliformis: tubo longissimo, gracili. Stigmata profundè incisa cristata.

16. CROCUS SATIVUS.

CHAR. SPEC. Stigmate tripartito longitudine corollæ reflexo, foliis linearibus margine revolutis.

ANGLICE, Saffron.

LOCUS. Helvetiæ, Pyrenæorum, Lusitaniæ, Thraciæ, Alpes. Perennis.

VIS. Subaromatica, luteo-tinctoria.

COMP. Syr. Croci. *Lond.* Tr. Croci Sativi. *Edin.* Conf. Aromatica. *Lond. Eblan.* Pil. Aloes cum Myrrh. *Lond.* Tr. Aloes Comp. *Lond. Edin. Eblan.* Tr. Cinch. Comp. *Lond. Eblan.* Tr. Aloes. Tr. Rhei. Comp. *Lond.*

IRIS. Willd. Spec. Plant. i, 224. Persoon Syn. i. 50.

ORDO NAT. ENSATÆ Linn. **IRIDES.** Juss.

CHAR. GEN. Corolla hexapetala, laciniis alternis reflexis. Stylus nullus. Stigmata petaliformia. Stamina incumbentia.

* Barbatae, foliis ensiformibus.

17. IRIS FLORENTINA.

CHAR. SPEC. Barbata, foliis ensiformibus, glabris, stapo subbifloro brevioribus, floribus sessilibus.

ANGLICE, Florentine Iris.

LOCUS. Rhodi Insula, Laconia. Perennis.

VIS. Fragrans. Inusitata.

CLASSIS III.

ORDO II.

TRIANDRIA. DIGYNIA.

SACCHARUM. Willd. Sp. Pl. i. 321. Persoon Syn. i. 102.

ORDO NAT. GRAMINA. Linn.

CLASSIS II. PLANTÆ MONOCOTYLEDONES
STAMINA HYPOGYNA. ORDO IV. GRAMINEÆ.
II. Styli duo. Stamina tria. Gluma uniflora. Juss.

CHAR. GEN. Flores omnes hermaphroditi. Calyx lanugine longa involucratus, bivalvis. Corolla uni-vel bivalvis, mutica, aut aristata. Stam. 1—3.

18. SACCHARUM OFFICINARUM.

CHAR. SPEC. Floribus paniculatis, geminatis, altero sessili, altero pedicellato. Corolla univalvi, mutica.

ANGLICE, Sugar-cane.

LOCUS. Indiæ Orientalis, et Occidentalis.

COMP. Syrupi omnes. *Lond. Edin. Eblan.* Trochisci omnes. *Edin. Conf. omnes. Lond.*

Inest huic plantæ saccharum, ut et plurimis aliis, variisque fructibus, Betæ maritimæ, Danco Carotæ, &c. et præsertim Aceri Saccharino, cujus succi evaporatione magnam sacchari copiam extrahunt Americani Septentrionales.

Valde mirum est quam abundè Saccharum Calcis solutionem in aquâ efficiat. Sacchari partes duæ Calcis unam aquâ priùs suspensam liquant.

*** Flores multiflori, vagi.

AVENA. Willd. Spec. Plant. i. 443. Persoon Syn. i. 100.

ORDO NAT. GRAMINA. Linn. CLASSIS II.

ORDO IV. GRAMINEÆ. VIII. Styli duo. Stamina tria. Glumæ multifloræ vagæ. Flores sæpius paniculati. Juss.

CHAR. GEN. Calyx bivalvis, multiflorus; arista dorsali, contorta.

19. AVENA SATIVA

CHAR. SPEC. Paniculata, calycibus dispermis, seminibus lævibus altero aristato.

ANGLICE, Common Oat.

LOCUS. Insula Juan Fernandez, Chili versus, ubi reperta est illustrissimo Anson, circiter annum 1741.

**** Spicati Receptaculo subulato.

SECALE. Willd. Spec. Plant. i. 471. Persoon Syn. i. 108.

ORDO NAT. GRAMINA. Linn. **CLASSIS II.**

ORDO IV. VII. Styli duo. Stamina tria. Glumæ multifloræ, supra axim seu rachim densè spicatæ.

CHAR. GEN. Calyx oppositus, bivalvis aut multiflorus, solitarius: glumis lineari lanceolatis lævibus aut utrinque canaliculatis.

20. SECALE CEREALE.

CHAR. SPEC. Glumarum ciliis scabris, squamis calycinis angustis minoribus. Spica passim ramosa.

ANGLICE, Rye.

LOCUS. Creta. Annua.

Secale cornutum, Calcar, Clavus secalinus, Secalis mater vel Secale luxurians, Gallicè ergot, et Anglicè, Cockspur of Rye. Cujus natura incerta est: auctoribus quibusdam vitium Secalis; aliis vero parasitus existimari videtur. Inter prioris sententiæ laudatores invenitur Johannes Bauhinus qui dicit "ut aristæ quædam postquam flores deciderunt; ac semina augmenta suscipere incipiunt, vitium aliquod in granis contrahunt. Tunc enim videmus quædam semina longius ex suis glumis protendi, et in mediocrem etiam crassitiem excrescere nigrumque colorem contrahere."

In Galliâ, et præsertim in provinciâ Sologne dictâ, Secale cornutum inter sana cerealis semina quotannis reperitur, æstatibusque pluvialibus quartæ fere messis parti adæquari

CLASSIS IV.

ORDO I.

TETRANDRIA. MONOGYNIA.

* *Flores Monopetali, Dicocci, Superi.*

Stellatæ.

RUBIA. Willd. Spec. Plant. i. 603. Persoon Syn. i. 129. Smith, Compend. Fl. Brit. 25.

ORDO NAT. STELLATÆ. Linn.

CLASSIS XI. DICOTYLEDONES MONOPETALÆ.

COROLLA EPIGYNA. ANTHERÆ DISTINCTÆ.

ORDO II. RUBIACEÆ. II. Fructus diccocus, dispermus.

Stamina sæpius quatuor. Folia plerumque verticillata; caulis plerumque herbaceus. Juss.

CHAR. GEN. Corolla monopetala campanulata quadrifida vel quinquefida. Baccæ duæ, monospermæ.

23. RUBIA TINCTORUM.

CHAR SPEC. Foliis lanceolatis, subsenis, annuis, caule aculeato scandente, Perennis.

ANGLICE, Madder.

LOCUS. Monspelii, in Italiæ, Helvetiæ, Danubii pratis.

VIS. Stiptica, rubro-tinctoria; urinam, et ossa tingens.

** Flores incompleti, inferi.

KRAMERIA. Willd. Spec. Plant. i. 693. Persoon Syn. i. 141.

ORDO NAT. INCERTÆ SEDIS. POLYPE-TALÆ GERMINE SUPERO. Juss.

CHAR. GEN. Calyx nullus. Corolla tetra-vel pentapetala. Nectarium tetraphyllum, biforme. Antheræ perforatæ. Drupa echinata.

24. KRAMERIA TRIANDRA.

CHAR. SPEC. Foliis oblongis, obovatisque, acuminatis, tomentosis, floribus triandris.

ANGLICE, Rhatany.

LOCUS. In Peruviae montium declivibus.

QUALITAS. Amara, Stiptica rubeo-tinctoria.

USUS. Febribus intermittentibus. Dysenteria, sed commodo dubio.

DOSIS. ʒi ad ʒi in pulvere.

DORSTENIA. Willd. Spec. Plant. i. 682. Persoon Syn. ii. 557.

ORDO NAT. SCABRIDÆ. Linn.

CLASSIS XV. ORDO III. URTICÆ. I. Flores in communi involucri monophyllo reconditi. Juss.

CHAR. GEN. Receptaculum commune monophyllum, carnosum in quo semina solitaria nidulantur.

25. DORSTENIA CONTRAJERVA.

CHAR. SPEC. Scapis radicatis, foliis pinnatifido-palmatis serratis, receptaculis quadrangulis.

ANGLICE, Contrajerva.

LOCUS. Peru, Mexico, Vera Crux, Vicentii Insula.

VIS. Aromatica, amara. Inusitata.

COMP. Pulvis Contrajervæ. Comp. Lond.

CLASSIS V.

ORDO I.

PENTANDRIA. MONOGYNIA.

* *Flores Monopetali, inferi, tetraspermi.*

Asperifoliæ.

ANCHUSA. Willd. Spec. Plant. i. 756. Persoon Syn. i. 159.

ORDO NAT. ASPERIFOLIÆ. Linn.

CLASSIS VII. ORDO IX. BORAGINÆÆ.

IV. Fructus gymno—tetraspermus. Faux corollæ instructa quinque squamis calcarum instar cavis, intra corollam prominulis in ejusdem laciniarum basi, extrâ hiulcis. Herbæ plerumque asperifoliæ. Juss.

CHAR. GEN. Corolla infundibuliformis; fauce clausa fornicibus. Semina basi insculpta, superficie ut plurimum venosa.

26. ANCHUSA TINCTORIA.

CHAR. SPEC. Tomentosa, foliis lanceolatis obtusis, staminibus corolla brevioribus.

ANGLICE, Alkanet.

LOCUS. Monspeli, et in Barbaria. Perennis.

VIS. Rubro-tinctoria.

* Flores Monopetali, inferi angiospermi.

MENYANTHES. Willd. Spec. Plant. i. 810. Persoon Syn. i. 171.

ORDO NAT. VAGÆ, ET INCERTÆ SEDIS. Linn.

CLASSIS VIII. ORDO I. LYSYMACHIÆ.

III. Genera Lysimachiis affinia. Juss.

CHAR. GEN. Corolla hirsuta. Stigma bifidum. Capsula unilocularis.

27. MENYANTHES TRIFOLIATA.

CHAR. SPEC. Corollis supernè dense barbatis, foliis ternatis.

ANGLICE, Buckbean.

LOCUS. In Europæ, et Americæ paludibus. Perennis.

VIS. Amarissima. Inusitata.

SPIGELIA. Willd. Spec. Plant. i. 824. Persoon Syn. i. 281.

ORDO NAT. STELLATÆ. Linn.

CLASSIS VIII. ORDO XIII. GENTIANÆ.

III. Capsula Didyma bilocularis. Juss.

CHAR. GEN. Calyx quinque partitus. Corolla infundibuliformis. Capsula didyma, bilocularis, tetra-valvis.

28. SPIGELIA MARILANDICA.

CHAR. SPEC. Perennis, caule tetragono, foliis omnibus oppositis.

ANGLICE, Perennial Wormgrass.

LOCUS. In America Boreali.

VIS. Amara, purgans. Inusitata.

CONVOLVULUS. Wild. Spec. Plant. i. 844. Persoon Syn. i. 177.

ORDO NAT. CAMPANACEI. Linn.

CLASSIS VIII. ORDO X. CONVULVULI.

I. Stylus unicus. Juss.

CHAR. GEN. Corolla campanulata, plicata. Stigmata duo, simplicia. Capsula bilocularis: loculis dispermis.

* Caule volubili.

29. CONVULVULUS SCAMMONIA.

CHAR. SPEC. Foliis sagittatis, postice truncatis, pedunculis teretibus subtrifloris.

ANGLICE, Scammony.

LOCUS. Syria, Mysia, Cappadocia. Perennis.

VIS. Acri-amaricans, casei odoris, purgans, drastica, anthelmintica, cito agens. Sæpenumero tormina inducit, ideoque aromaticis commisceri oportet.

DOSIS. Gr. v ad xv.

COMP. Conf. Scammoniae. Lond. Eblan. Pulv. Scammon. Comp. Lond. Edin. Ext. Colocynth. Comp. Lond. Pulv. Sennæ Comp. Lond.

30. CONVULVULUS JALAPA. Willd. Spec. Plant. i. 860. Persoon Syn. i. 179.

CHAR. SPEC. Caule volubili, foliis ovatis, subcordatis, obtusis, obsolete repandis subtus villosis, pedunculis unifloris.

ANGLICE, Jalap.

LOCUS. Mexico, Vera Cruz. Zalappæ oppido nuncupatus.

VIS. Nauseosa, acris: inter omnia purgantia, quæ hodierno in usu sunt, nullum Jalapâ præstantius; quia viarum primarum evacuationem celeritèr et penitèr efficit, præsertim Calomelane addito; et cum potassæ supertartrate mixta copiosas dejectiones fluidas excitat.

DOSIS. Gr. v ad ʒi.

DATURA. Willd. Spec. Plant. i. 1007. Persoon Syn. i. 216.

ORDO NAT. LURIDÆ. Linn.

CLASSIS VIII. ORDO VIII. SOLANÆÆ. I. Fructus capsularis. Juss.

CHAR. GEN. Corolla infundibuliformis, plicata. Calyx tubulosus, angulatus, deciduus; basi orbiculata persistente. Capsula tetravalvis, tetralocularis, lævis aut spinosa.

31. DATURA STAMONIUM.

CHAR. SPEC. Pericarpis spinosis erectis ovatis, foliis ovatis sinuatis glabris basi in petiolum attenuatis.

ANGLICE, Thornapple.

LOCUS. America. Nunc vulgaris est planta per Europam. Perennis.

VIS. Stimulans, narcotica; sed viribus incertis, et pluribus aliis posthabenda. Foliorum siccatorum fumus, Asthmâ spasmodicâ laborantes, inhalatus aliquando, Tabaci more, relevat. Dosibus nimis grandioribus exhibita vomitum, vertiginem, delirium, stuporem, convulsiones et alia molesta ciet. Remedia, emetica, stimulantia.

D. Brandes nuper ostendit Stramonii vires de materiâ novâ alkalinâ naturâ Daturiam dictam pendere.

ERYTHRÆA. Willd. Spec. Plant. i. 1065. Persoon Syn. i. 283.

ORDO NAT. ROTACÆÆ. Linn.

CLASSIS VIII. ORDO XIII. GENTIANÆ.

II. Capsula simplex bilocularis. Juss.

CHAR. GEN. Calyx appressus, subpentagonus, quinque-dentatus. Corolla infundibuliformis: tubo longo. Antheræ, post anthesin, spiraliter tortæ. Stigmata duo approximata, rarius simplex. Capsula sublinearis, unilocularis. Receptacula duo, extantia.

* Floribus sessilibus.

32. ~~34~~ ERYTHRÆA CENTAURIUM.

CHAR. SPEC. Foliis oblongo-ovalibus basi rosaceis congestis, caule elongato, floribus fasciculato-paniculatis, basi squamis unicâ vel duabus cinctis.

GENTIANA CENTAURIUM. Linn.

CHIRONIA CENTAURIUM. Willd.

ANGLICE, Common Centaury.

LOCUS. Angliæ indigena.

VIS. Tota herba amara. Inusitata.

33. ~~32~~ HYOSCYAMUS. Willd. Spec. Plant. i: 1010. Persoon Syn. i. 217.

ORDO NAT. LURIDÆ. Linn.

CLASSIS VIII. ORDO VIII. SOLANÆÆ. I. Fructus Capsularis. Juss.

CHAR. GEN. Corolla infundibuliformis, obtusa, irregularis. Stamina inclinata. Capsula operculata, bilocularis.

HYOSCYAMUS NIGER.

CHAR. SPEC. Foliis amplexicaulibus, sinuatis, floribus venosis sessilibus.

ANGLICE, Henbane.

LOCUS. Europæ ruderatis pinguibus. Biennis sive annua.

VIS. Mucilaginosa, virosa.

COMP. Tinct. Hyoscyami. Ext. Hyoscyami. *Lond. Edin. Eblan.*

USUS. Recentes medici hunc anodynum opio subinde anteponendum putaverunt, quia alvi obstipationem non

efficit, sed potius vim laxativam possidet. Nihilominus nauseam, vertiginem, &c. aliquando excitat. Belladonnæ more, pupillæ oculi fines extendit, nec tamen viribus tam certis aut fortibus. Docet D. Brandes huic plantæ inesse alkali proprium Hyoscyomiam dictum.

NICOTIANA. Willd. Spec. Plant. i. 1014. Persoon Syn. i. 217.

ORDO NAT. LURIDÆ. Linn.

CLASSIS VIII. ORDO VIII. SOLANÆÆ. Juss.

CHAR. GEN. Corolla infundibuliformis, limbo plicato. Stamina inclinata. Capsula bivalvis, bilocularis.

34. ~~33.~~ NICOTIANA TABACUM.

CHAR. SPEC. Foliis lanceolato-ovatis sessilibus decurrentibus, floribus acutis.

ANGLICE, Tobacco.

LOCUS. America. Annuæ.

VIS. Nauseosa, acris, virosa, anodyna, narcotica, vomitoria, errhina.

COMP. Infusum Tabaci. Lond. Vinum Nicotianæ Tabaci. Edin.

Vires sedativæ potentissimæ, et auctore D. Brodie, cordis actionem diminuendo, per systema nervosum,ducuntur.

Herniis et urinæ retentione laborantibus, spasmus subindè feliciter laxat, sed quam cautissime cavendum est, ne syn-copen lethalem inducat. Contra talem casum, Ammonia et alia stimulantia confestim adhibenda.

DOSIS. Semidrachmam vel certe drachmam, in aquæ ferventis octario nequaquam excedat, enematis formâ.

Olei essentialis e foliis destillati, guttula unica linguæ super-fusa canem protinus interfecisse fertur.

* Fructus baccatus aut drupaceus.

STRYCHNOS. Willd. Spec. Plant. i. 1052. Persoon Syn. i. 264.

ORDO NAT. CLASSIS VIII. ORDO XIV. APOCINEÆ. IV. Genera Apocineis affinia, non lactescentia. Juss.

CHAR. GEN. Corolla tubulosa, quinquifida. Bacca unilocularis, cortice lignoso.

35. STRYCHNOS NUX VOMICA.

CHAR. SPEC. Foliis ovatis, caule inermi.

VULGO. Nux vomica.

LOCUS. Malabaria, Zeylona. Perennis.

QUAL. Amara, virosa.

Anno 1818, D. Pelletier et Caventou materiem alkalinam in hujus plantæ fructu detexere, cui nomen Strychniam tribuere. Alkohole fervente facile liquatur, sed partes aquæ frigidaë 6000 Strychniæ unicam modo solvunt: et solutio quamvis tenuissima gustum amarissimum præstat. Toxicorum omnium vegetabilium hoc penè valentissimum, et admiratione dignum est; namque grani dimidium cuniculi gutturi inflatum horæ duodecimo spatio mortem efficit, trismo et tetano inductis.

CAPSICUM, (καπτω, mordeo.) Willd. Spec. Plant. i. 1050. Persoon Syn. i. 229.

ORDO NAT. LURIDÆ. Linn.

CLASSIS VIII. ORDO VIII. SOLANEÆ. II. Fructus baccatus. Juss.

CHAR. GEN. Corolla rotata, antheræ conniventes, bacca bilocularis, exsucca.

36. CAPSICUM ANNUUM.

CHAR. SPEC. Caule herbaceo, pedunculis solitariis.

ANGLICE, Cayenne Pepper.

LOCUS. Brazilia, Mexico, Barbados. Annua.

QUAL. Acris, stimulans.

USUS. In Dyspepsia, et Tonsillarum ulceribus.

DOSIS. Gr. i ad ij pilulari formâ.

COMP. Tinctura Capsici. *Lond. et Eblan.*

SOLANUM. Willd. Spec. Plant. i. 1025. Persoon Syn. i. 220.

ORDO NAT. LURIDÆ. Linn.

CLASSIS VIII. ORDO VIII. II. Fructus baccatus.

SOLANÆÆ. Juss.

CHAR. GEN. Calyx persistens. Corolla rotata, aut campanulata quinqueloba, plicata. Antheræ subcoalitæ, apice poro gemino, dehistentes. Bacca bilocularis, polysperma.

* *Inermia* foliis integris, aut sinuatis.

37. SOLANUM DULCAMARA.

CHAR. SPEC. Caule inermi fruticoso scandente, foliis cordatis glabris superioribus auriculatis, corymbis oppositifoliis.

ANGLICE, Bitter-sweet.

LOCUS. Europæ sepes humidiusculæ. Perennis.

QUAL. Nauseosa, dulcamara.

USUS. In morbis cutaneis.

COMP. Dec. Dulcamaræ. *Lond.*

Dulcamaræ buccæ purgantes et emeticæ sunt. Cani datæ intra trium horarum spatium eum occiderunt.

ATROPA. ATROPOS.

(*a τρεπω*, verto,) Furiarum ultima quæ averti nequeat, et vitæ filum findit. Willd. Spec. Plant. i. 1016. Persoon Syn. i. 218.

ORDO NAT. LURIDÆ. Linn.

CLASSIS VIII. ORDO VIII. SOLANÆÆ. Juss.

CHAR. GEN. Corolla campanulata. Stamina distantia. Bacca globosa, bilocularis, calyci insidens.

38. ATROPA BELLADONNA.

CHAR. SPEC. Caule herbaceo brachiato, foliis ovatis integris.

ANGLICE, Deadly Nightshade.

LOCUS. Austriæ, Angliæ, Germaniæ, Italiæ montibus sylvos. Perennis.

QUALITAS. Narcotica, anodyna. Amara, valde venenata. Ita sopit vim irritabilem ventriculi, secundum Hallerum, ut vix quatuordecim grana Antimonii Tartrisi a sumptis baccis vomitum excitarunt.

USUS. Plurimis morbis intrinsicis adhibita fuit Belladonna: dolorem subindè levat, veruntamen delirium, sopor convulsiones, visus diminutus et alia periculosa usum ejus aliquando sequuntur. Quibus perpensis, medicus prudens remedium tam incertum, quasi anodynum, in praxi rarò tentabit. Sed fatendum est Belladonnam extrinsecus applicatam multum prodesse in Rheumatismo, Dysmenorrhæa; et præsertim, pupillæ fines extendendo, in Iritide, et oculi suffusione, (Anglice Cataract.) Auctore, et suasore, D. Himly de Göttingen.

Continet propriam materiem, naturâ alkalinâ, Atropiam dictam, D. Brandes nuper patefactam.

COMP. Ext. Belladonnæ. *Lond.* Dosis, gr. $\frac{1}{4}$ ad gr. i. Succus spissatus Atropæ Belladonnæ. *Edin.*

Flores monopetali superi.

CINCHONA. Willd. Spec. Plant. i. 957. Persoon Syn. i. 196.

ORDO NAT. CYMOSÆ. Linn.

CLASSIS XI. ORDO II. RUBIACEÆ. IV. Fructus mono-carpus bilocularis polyspermus. Stamina quinque. Folia opposita; caulis sæpe frutescens. Juss.

CHAR. GEN. Calyx campanulatus, quinque-dentatus. Corolla infundibuliformis: limbo quinque partito ut plurimum hirto. Capsula oblongo-teres, bilocularis, bipartibilis: dissepimento parallelo. Semina numerosa, compressa margine membranaceo cincta.

* Corollis extus sericeo-tomentosis, limbo suprâ barbatis.

CINCHONA CONDAMINEA. Plantæ Æquinoxiales, tom. i. 33. tab. 10.

39. CINCHONA LANCEOLATA. Flora Peruviana, tom. ii. 51. t. 223. Et ejusdem operis, tom. ii. tab. 191.

CINCHONA NITIDA.

CHAR. SPEC. Foliis ovato-lanceolatis utrinque ramulisque glabris, dentibus calycinis ovatis acuminatis stigmate emarginato, capsulis ovatis costatis. D. Don. Mss.

ANGLICE, Quill Bark.

LOCUS. America australior.

40. CINCHONA CORDIFOLIA.

CHAR. SPEC. Foliis cordatis ellipticisve subtus ramulisque pilosiusculis, dentibus calycinis rotundatis, stigmate bilobo, capsulis ellipticis lævibus.

ANGLICE, Yellow Bark. D. Don. Mss.

**** Corollis extus sericeo-tomentosis, limbo suprà glabris.**

41. CINCHONA MAGNIFOLIA. Flora Peruviana, tom. ii. 53, t. 196.

CHAR. SPEC. Foliis ellipticis subtus ramulisque tomentosis, dentibus calycinis ovatis acutis, stigmate exserto bipartito, capsulis cylindricis. D. Don. Mss.

ANGLICE, Red Bark.

VIS. Cinchona Condaminea morborum curatione specierum omnium valentissima. Ante annum 1818 parum de vera ejus natura apud Physicos compertum fuit; quando DD. Pelletier et Caventou proprium alkali vegetabile quod in crystallos abire solet, in ejus Cortice detexere, cui Cinchoniam nomen dedere. Hoc alkali cum acido Kinico nuncupato conjunctum reperitur, et de sale ita constituto medicamenti vires pendent.

Cinchona Cordifolia, Prioris speciei vim medicinalem haud adæquat. DD. Pelletier et Caventou in ejus cortice alium alkali invenere Quiniam dictum. Quinia crystallinam formam nunquam induit, attamen acido Kinico consociata Quiniæ Super-Kinatem format.

Cinchona magnifolia haud æquè amara sed tamen vi as-

tringente Cinchonam Cordifoliam superat; Cinchonæ et Quinæ Super-Kinatam parvâ quantitate continet.

Quinia, et Cinchonia acidis variis consociatæ, aliorum alkalium vegetabilium more, sales proprios conficiunt. Quinæ Sulphatem, in hodierna praxi, mazzere Cinchonæ corticis pene fungi, omnibus satis cognitum est.

Ad Quinæ Sulphatem obtinendam, acido Sulphurico aquâ valdè diluto Cinchonæ Cordifoliæ corticem contusum adjice, coque per horam, et cola. Liquori colato calcem adde. Pulverem ita demissum exsicca, et in Alkohole macera, donec omnis Quinia soluta sit. Alkohol destilla, et residuæ Quinæ acidum sulphuricum dilutum, ad plenam saturationem adde, solve, et aquam consume ut in crystallos prodeat.

Omnium Medicaminum tonicorum Cinchona facile princeps. In Europam anno 1640, propter vires in Febrium Intermittentium curatione, primùm invehebatur: earum tamen parexysmo calido medicamenti dosin imminui oportet. Febribus remittentibus, et aliis perutile est hoc remedium; at potissimum notandum est, nihil prodesse illis affectionibus de urethræ stricturâ pendentibus, et febrium speciem simulantibus. Tam frequenter nuperrimè adhiberi solet, ut difficile esset omnes morbos hoc loco indicare quibus optimè conveniat. Præcipuè autem datur pro Rheumatismo, Hemigranæa; Cynanche, Rubeolâ, Scarlatinâ, et Variolâ malignis, et multis aliis, naturæ intermittētis, morbis.

DOSIS. Pulveris corticis gr. x. ad ʒi. Decocti vel Infusi ʒi ad ʒij. Quinæ Sulphatis gr. ss. ad gr. x. solutionis, vel pilulari formâ.

COMP. Infusum. *Lond. Edin. Eblan.* Decoctum. *Lond. Edin. Eblan.* Extractum Cinchonæ. *Lond. Edin.* Extractum Cinchonæ Resinosum. *Lond. Eblan.* Tinctura. *Lond. Edin. Eblan.* Tinctura Cinchonæ Comp. *Lond. Edin. Eblan.*

42. **CALLICOCCA IPECACUANHA.** Brotero Act. Societat. Linn. *Lond.* tom. vi. p. 137.

Flores aggregati in capitulum solitarium; Calyx quinque-dentatus. Corolla monopetala: tubus cylindræus,

longua, suprà parùm ampliatus, fauce et extùs lanuginosus; Stigmata duo, oblonga, crassiuscula, obtusa, antherarum longitudine. Bacca unilocularis, disperma. Caulis suffruticosus.

Folia inferiora caduca, ita ut in plantæ florescentia, 4, 6, vel 8 solidum, rarissimè plura, ad apicem caulis persistent.

ANGLICE, *Ipecacuban*.

LOCUS. In solo umbroso, humerosoque sylvarum, quæ Pernambucoiâ, Bahiâ, Rio Janeiriâ, aliisque Brasiliæ provinciis abundant. Perennis.

VIS. Nauseosa, amara, pulvis subtilis ære suspensus, interdum pulmones valdè molestat.

USUS. Dosibus potentioribus hæc omnibus emeticis vegetabilibus præstat, quia efficacitè et lenitè agit; symptomata febrilia, morbo incipiente mitigat; vomitum, nauseam diurnam, sudorem, diuresin et catharsin subinde excitando; præcipuè si Antimonii Tartarizati granum ad Ipecacuanhæ gr. xv. vel scrupulum adjiciatur.

Dosibus minoribus utilissima est in Diarrhœa et Dysenteria: ex intervallis horarum trium quatuorve sumpta, Ipecacuhana diaphoretica, et expectorans est in Pneumonia, Pleuritide, et aliis Phlegmaciis. Reperiere DD. Majendie et Pelletier, materiem pollentem ab Ipecacuanhâ secerni posse: hanc materiem Emetinam nominaverunt.

DOSIS. Ad Vomitus ciendum gr. v. ad ʒi: ad Diaphoresin, et Expectorationem excitandam grani quadrans ad grana duo.

COMP. Pulvis Ipecac. Comp. *Lond. Eblan.*

Vinum. Ipecacuanhæ. *Lond. Edin. Eblan.*

* Flores pentapetali inferi.

RHAMNUS. Willd. Spec. Plant. i. 1092. Persoon Syn. i. 238.

ORDO NAT. VEPRECULE. Linn.

CLASSIS XIV. PLANTÆ DICOTYLEDONES
POLYPETALÆ STAMINA PERIGYNA.

ORDO XIII. RHAMNI. III. Stamina petalis opposita. Fructus drupaceus. Juss.

CHAR. GEN. Calyx campanulatus, 4-5-fidus. Corollæ squamæ stamina munientes, calyci insertæ. Stigma 1-2-4-fidum, Bacca 3-4-sperma. (Flores passim polygami, et dioici.)

* Spinosi.

43. RHAMNUS CATHARTICUS.

CHAR. SPEC. Spinis terminalibus, floribus quadrifidis dioicis, foliis ovatis serratis. Stigma quadrifidum.

ANGLICE, Buckthorn.

LOCUS. Europæ saltus. Perennis.

VIS. Amara, luteo-tinctoria, purgans, drastica.

COMP. Syr. Rhamni, *Lond. Edin.*

VITIS. Willd. Spec. Plant. p. 1180. Persoon Syn. i. 252.

ORDO NAT. SARMENTACEÆ. Linn.

CLASSIS XIII. PLANTÆ DICOTYLEDONES.

POLYPETALÆ. STAMINA HYPOGYNA.

ORDO XII. VITES. Juss.

CHAR. GEN. Petala apice calyptratim cohærentia, emarcida. Bacca pentasperma, supera.

44. VITIS VINIFERA.

CHAR. SPEC. Foliis lobatis sinuatis nudis aut tomentosis.

ANGLICE, Vine.

LOCUS. Orbis totus temperatus. Perennis.

VIS. Acido-dulcis. Fructus antisepticus.

DIOSMA. Willd. Spec. Plant. i. 1133. Persoon Syn. i. 246.

CLASSIS XIII. ORDO XXI. RUTACEÆ.

III. Genera Rutaceis affinia. Juss.

CHAR. GEN. Calyx sub-quinque-partitus persistens. Corolla pentapetala. Nectaria quinque, supra germen. Capsulæ tres sive quinque, coalitæ. Semina Calyptrata.

45. DIOSMA CRENATA.

CHAR. SPEC. Foliis ovatis crenatis subtus punctatis, floribus axillaribus solitariis. Calycis folia subulata, crenata.

ANGLICE, Buchu.

LOCUS. Promontorium Bonæ Spei. Perennis.

VIS. Aromatica, sudorifica, diuretica.

VIOLA. Willd. Spec. Plant. i. 1159. Persoon Syn. i. 253.

ORDO NAT. CAMPANACEI. Linn.

CLASSIS XIII. ORDO XX. CISTI. Genera Cistis affinia fructu trivalvi valvis seminiferis, sed definitè staminifera. Juss.

CHAR. GEN. Calyx pentaphyllus. Corolla pentapetala, irregularis, posticè cornuta (aut ecalcerata.) Anthere apice membranula cohærentes, aut distinctæ. Capsula supera, trivalvis, unilocularis.

Acaules.

46. VIOLA ODERATA.

CHAR. SPEC. Foliis cordatis, stolonibus reptantibus.

ANGLICE, Sweet Violet.

LOCUS. Europæ nemoribus, et ad sepes.

VIS. Ceruleo-tinctoria.

COMP. Syr. Violæ. *Edin. Eblan.*

CLASSIS V.**ORDO II.**

PENTANDRIA. DIGYNIA.

Flores Monopetali inferi.

GENTIANA. Willd. Spec. Plant. i. 1331. Persoon Syn. i. 284.

ORDO NAT. ROTACEÆ. Linn.

CLASSIS VIII. ORDO XIII. GENTIANÆ. Capsula simplex unilocularis. Juss.

CHAR. GEN. Corolla monopetala. Capsula bivalvis, unilocularis. Receptaculis duobus, longitudinalibus.

* Corollis 5—9 fidis subcampanulatis.

47. GENTIANA LUTEA.

CHAR. SPEC. Foliis latis ovatis nervosis, corollis subquinqfidis rotatis verticillatis. Calycibus spathaceis.

ANGLICE, Gentian.

LOCUS. In alpinis et subalpinis pratis. Perennis.

VIS. Amarissima. Tonica præstans, parum nauseosa.

USUS. In Dyspepsiâ.

COMP. Ext. Gentianæ. *Lond. Edin. Eblan.* Inf. Gent. Comp. *Lond. Edin. Eblan.* Tinct. Gent. Comp. *Lond. Edin. Eblan.* Vinum. Gent. Comp. *Edin.*

Flores Incompleti.

ULMUS. Willd. Sp. Pl. i. 1324. Persoon Syn. i. 290.

ORDO NAT. STATUMINATÆ. Linn.

CLASSIS XV. ORDO IV. AMENTACEÆ. I. Flores. Hermaphroditi. Juss.

CHAR. GEN. Calyx campanulatus quinqfidus. Corolla nulla. Capsula compressa, membranaceo-alata. (Stamina etiam quatuor vel octo. Flores conglomerati.)

48. ULMUS CAMPESTRIS.

CHAR. SPEC. Foliis duplicato-serratis basi inæqualibus, floribus subsessilibus conglomeratis, pentandris, fructibus glabris.

ANGLICE, Elm.

LOCUS. In campestribus. Perennis.

VIS. Subamara, mucilaginosâ.

COMP. Dec. Ulmi. *Lond. Eblan.*

Flores Pentapetali superi dispermi. Umbellatæ.

A. Involucro universali, partialique.

ERYNGIUM. Willd. Spec. Plant. i. 1356. Persoon Syn. i. 299.

ORDO NAT. UMBELLATÆ. Linn.

**CLASSIS XII. PLANTÆ DICOTYLEDONES
POLYPETALÆ. STAMINA EPIGYNA.**

ORDO II. UMBELLIFERÆ. IV. Umbelliferæ anomalæ. Juss.

CHAR. GEN. Flores capitati. Involucrum polyphyllum. Calyx proprius, pentaphyllus, superus. Corolla pentapetala. Receptaculum paleaceum. Fructus coronatus, bipartibilis.

* Foliis laciniatis.

49. ERYNGIUM MARITIMUM.

CHAR. SPEC. Foliis radicalibus subrotundis plicatis spinosis, capitulis pedunculatis, paleis tricuspidatis.

ANGLICE, Eryngo.

LOCUS. In maritimis. Perennis.

VIS. Diuretica, aperiens, existimatur.

DAUCUS. Willd. Spec. Plant. i. 1389. Persoon Syn. i. 306.

ORDO NAT. UMBELLATÆ. Linn.

CLASSIS XII. ORDO II. UMBELLIFERÆ.

III. Umbelliferæ Veræ.

Umbellæ et Umbellulæ Involucratæ. Juss.

CHAR. GEN. Corollæ subradiatæ. Flosculi disci abortivi. Fructus pilis hispidus. (Umbella fructus contractus infundibuliformis.)

50. DAUCUS CAROTA.

CHAR. SPEC. Seminibus hispidis, petiolis subtus nervosis, foliorum laciniis angusto-linearibus acutis.

ANGLICE, Carrot.

LOCUS. In pascuis, arvis. Biennis.

USUS. Cataplasmata e dauci radice confecta, ulceribus fœdis utilia, putorem sæpe corrigendo.

CONIUM. Willd. Spec. Plant. i. 1395. Persoon Syn. i. 308.

ORDO NAT. UMBELLATÆ. Linn.

CLASSIS XII. ORDO II. UMBELLIFERÆ. Juss.

CHAR. GEN. Involucella dimidiata, subtriphylla.

Fructus subglobosus, quinque-striatus, utrinque crenatus.

51. CONIUM MACULATUM.

CHAR. SPEC. Caule ramosissimo maculato, seminibus striatis.

ANGLICE, Hemlock.

LOCUS. In agris, ruderatis. Biennis.

VIS. Amara. Nauseosa. Narcotica.

USUS. Loco Opii Conium sæpe suppositum est ; quoniam niam dolorem mitigat, sed neque systema vasculosum concitat, nec alvi astrictionem efficit. In Rheumatismo, et Pertussi multum valet, sed adhibenda est cura ne modus idoneus transeat, nam vertigo, convulsiones, et talia molesta deinde sequuntur.

Cataplasmatis formâ hæc planta contusa carcinomatibus ulcerosis, et dolorosis interdum applicatur.

COMP. Ext. Conii. *Lond. Edin. Eblan.*

DOSIS. Pulveris, et Extracti Conii gr. iii. ad ʒi.

SIUM. Willd. Spec. Plant. i. 1431. Persoon Syn. i. 316

ORDO NAT. UMBELLATÆ. Linn.

CLASSIS XII. ORDO II. UMBELLIFERÆ. Juss.

CHAR. GEN. Fructus subovatus, compressus, striatus.

Involucrum polyphyllum. Petala cordata.

52. SIUM NODIFLORUM,

CHAR. SPEC. Caule procumbente, foliis pinnatis foliis ovatis, umbellis axillaribus, sessilibus. Involucrum sæpè deest, aut monophyllum.

ANGLICE, Procumbent Water Parsnip.

LOCUS. Ad ripas fluviorum. Perennis.

VIS. Antiscorbuticum olim existimatum. Inusitatum.

CUMINUM. Willd. Spec. Plant. i. 1440. Persoon Syn. i. 318.

ORDO NAT. UMBELLATÆ. Linn.

CLASSIS XII. ORDO II. UMBELLIFERÆ. Juss.

CHAR. GEN. Fructus ovatus, striatus. Umbellulæ quatuor. Involucra quadrifida. Foliis simplicibus, et trifidis.

53. CUMINUM CYMINUM.

ANGLICE, Cumin.

LOCUS. Æthiopia, Ægyptus. *Annua.*

VIS. Acris aromatica.

COMP. Emp. Camini. *Lond.*

FERULA. Willd. Spec. Plant. i. 1411. Persoon Syn. i. 311.

ORDO NAT. UMBELLATÆ. Linn.

CLASSIS XII. ORDO II. UMBELLIFERÆ. Juss.

CHAR. GEN. Involucrum universale caducum; parziale polyphyllum. Fructus ovatus, compresso-planus, striis utrinque tribus.

54. FERULA ASAFŒTIDA.

CHAR SPEC. Foliolis alternatim sinuatis obtusis.

ANGLICE, Asafœtida.

LOCUS. Persia. *Perennis.*

VIS. Odore alliaceo, foetido. Carminativa. Antispasmodica?

USUS. In Hysteriâ, Hypochondriasi, Dyspepsiâ: beneficio ancipiti.

COMP. Mist. Asafœtidæ *Lond. Eblan.* Tinct. Asafœtidæ. *Lond. Edin. Eblan.* Sp. Ammon. Foetidus. *Lond. Eblan.* Tinct. Castorei Comp. *Eblan.* Pil. Asafœtidæ Comp. *Edin.* Pil. Aloes et Asafœtidæ. *Eblan.* Pil. Galb. Comp. *Lond.* Enema Foetidum. *Eblan.*

DOSIS. Gr. v. ad ʒi.

BUBON. Willd. Spec. Plant. i. 1439. Persoon Syn. i. 317.

ORDO NAT. UMBELLATÆ. Linn.

CLASSIS XII. ORDO II. UMBELLIFERÆ. Juss.

CHAR. GEN. Involucra polyphylla. Fructus ovatus, striatus, villosus et glaber.

**** Fruticosa.**

55. BUBON GALBANUM.

CHAR. SPEC. Foliolis ovato cuneiformibus acutis argute serratis, umbellis paucis, seminibus glabris caule frutescente glauco.

ANGLICE, Galbanum.

LOCUS. Æthiopia. Perennis.

VIS. Resina graveolens, calida.

USUS. Hysteria.

COMP. Pil. Galb. Comp. *Lond.* Pil. Asafœtidæ Comp. *Edin.* Pil. Myrrhæ Comp. *Eblan.* Tinct. Galb. *Eblan.* Emp. Galb. *Eblan.* Emp. Galb. Comp. *Lond.* Emp. Asafœtidæ. *Edin.* Emp. Gummosum. *Edin.*

DOSIS. Gr. x. ad ʒi.

ANGELICA. Willd. Spec. Plant. i. 1428. Persoon Syn. i. 315.

ORDO NAT. UMBELLATÆ. Linn.

CLASSIS XII. ORDO II. UMBELLIFERÆ. Juss.

CHAR. GEN. Fructus subrotundus, solidus, utrinque tri-alatus. Stylis reflexis. Corollæ æquales; petalis incurvis.

56. ANGELICA ARCHANGELICA.

CHAR. SPEC. Foliorum impari lobato.

ANGLICE, Garden Angelica.

LOCUS. Ad rivulos. Perennis.

VIS. Acris, amara.

B. Involucris partialibus.

CORIANDRUM. Willd. Spec. Plant. i. 1448. Persoon Syn. i. 319.

ORDO NAT. UMBELLATÆ. Linn.

CLASSIS XII. ORDO II. UMBELLIFERÆ. Juss.

CHAR. GEN. Corolla radiata. Petala inflexo-emar-

ginata. Involucrum universale, monophyllum; partialia dimidiata. Fructus sphaericus.

57. CORIANDRUM SATIVUM.

CHAR. SPEC. Fructibus globosis. Calyces stylique persistentes.

ANGLICE, Coriander.

LOCUS. Europæ agri. Annuæ.

VIS. Carminativa.

COMP. Conf. Sennæ. *Lond.*

C. Involucro nullo, nec universali, nec partialibus.

CARUM. Willd. Spec. Plant. i. 1470. Persoon Syn. i. 323.

ORDO NAT. UMBELLATÆ. Linn.

CLASSIS XII. ORDO II. UMBELLIFERÆ. Juss.

CHAR. GEN. Fructus ovato-oblongus, striatus. Involucrum sub-monophyllum. Petala carinata, inflexo-emarginata.

58. CARUM CARUI.

CHAR. SPEC. Caule ramoso, vaginis foliorum ventricosis, involucro partiali nullo.

ANGLICE, Carraway.

LOCUS. Europæ prata. Biennis.

VIS. Acris, aromatica.

COMP. Oleum Carui. *Lond. Eblan.* Aqua Carui. *Lond.* Sp. Carui. *Lond. Edin. Eblan.*

PASTINACA. Willd. Spec. Plant. i. 1465. Persoon Syn. i. 322.

ORDO NAT. UMBELLATÆ. Linn.

CLASSIS XII. ORDO II. UMBELLIFERÆ. Juss.

CHAR. GEN. Fructus ellipticus, compresso-planus. Petala involuta, integra.

59. PASTINACA OPOPONAX.

CHAR SPEC. Foliis pinnatis. Foliolis basi antica excisis. Petioli subhispidi.

ANGLICE, Opoponax.

LOCUS. Italia, et Sicilia. Perennis.

VIS. Amara, acris, graveolens.

ANETHUM. Willd. Spec. Plant. i. 1469. Persoon
Syn. i. 323.

ORDO NAT. UMBELLATÆ. Linn.

CLASSIS XII. ORDO II. UMBELLIFERÆ. Juss.

CHAR. GEN. Fructus subovatus, compressus, striatus.

Petala involuta, integra, (flava.)

60. ANETHUM GRAVEOLENS.

CHAR. SPEC. Annuum, fructibus compressis.

ANGLICE, Dill.

LOCUS. Hispania, Lusitania.

VIS. Aromatica, Carminativa.

USUS. Colicâ infantum.

COMP. Aqua Anethi. *Lond.*

61. ANETHUM FœNICULUM.

CHAR. SPEC. Perenne, fructibus ovatis.

ANGLICE, Fennel.

LOCUS. Gallia.

VIS. Aromatica, Carminativa.

COMP. Aqua Fœniculi. *Lond. Eblan.* Oleum Sem :
Fœniculi dulcis. *Eblan.*

PIMPINELLA. Willd. Spec. Plant. i. 1471. Per-
soon Syn. i. 323.

ORDO NAT. UMBELLATÆ. Linn.

CLASSIS XII. ORDO II. UMBELLIFERÆ. Juss.

CHAR. GEN. Fructus ovato-oblongus. Petala inflexa,
Stigmata subglobosa. Umbellulæ subglobosæ, primo nu-
tantes.

62. PIMPINELLA ANISUM.

CHAR. SPEC. Foliis radicalibus trifidis incisis. In-
volucrum monophyllum. Fructus pubescens.

ANGLICE, Anise seeds.

LOCUS. Syria, Ægyptus. Annua.

VIS. Aromatica.

COMP. Oleum Anisi. *Lond. Edin. Eblan.* Sp. Anisi. *Lond.*

CLASSIS V.

ORDO III.

PENTANDRIA. TRIGYNIA.

* *Flores Superi.*

SAMBUCUS. Willd. Spec. Plant. i. 1494. Persoon Syn. i. 328.

ORDO NAT. DUMOSÆ. Linn.

CLASSIS XI. DICOTYLEDONES MONOPETALÆ.
COROLLA EPIGYNA. ANTHERÆ DISTINCTÆ.

ORDO III. CAPRIFOLIA. III. Calyx¹ bracteatus.
Stylus nullus; stigmata tria. Corolla monopetala. Juss.

CHAR. GEN. Calyx quinquepartitus. Corolla quinquefida. Bacca trisperma.

63. SUMBUCUS NIGRA.

CHAR. SPEC. Cymis quinque partitis, stipulis duabus utrinque setaceis, foliis pinnatis, foliolis ovatis serratis, caule arboreo.

ANGLICE, Elder.

LOCUS. Germania, Japonia. Perennis.

VIS. Acris, nauseosa.

COMP. Sucus spissatus Sambuci nigri. *Edin. Eblan.*
Ung. Sambuci. *Lond. Eblan.*

* *Flores inferi.*

RHUS. Willd. Spec. Plant. i. 1477. Persoon Syn. i. 324.

ORDO NAT. DUMOSÆ. Linn.

CLASSIS XIV. PLANTÆ DICOTYLEDONES
POLYPETALÆ. STAMINA PERIGYNA. ORDO
XII. TEREBINACEÆ. I. Germen simplex. Fructus
unilocularis monospermus. Juss.

CHAR. GEN. Calyx quinque partitus. Petala quin-
que. Bacca monosperma.

** Foliis ternatis.

64. RHUS TOXICODENDRON.

CHAR. SPEC. Foliolis petiolatis angulatis pubescentibus, caule radicante.

ANGLICE, Poison Oak.

LOCUS. America Septentrionalis. Perennis.

VIS. Stimulans, Narcotica.

USUS. In Paralyti, sed parum valet.

DOSIS. Gr. i. ad v.

CLASSIS V.

ORDO V.

PENTANDRIA. PENTAGYNIA

* *Flores inferi.*

LINUM. Willd. Spec. Plant. i. 1533. Persoon Syn.
i. 334.

ORDO NAT. SUCCULENTÆ. Linn.

CLASSIS XIII. ORDO XXII. CARYOPHYLLÆ.

VII. Genera Caryophylleis affinia. Juss.

CHAR. GEN. Calyx pentaphyllus. Petala quinque.
Capsula quinque-valvis, decem-locularis. Semina solitaria.
(Filamenta basi coalita.)

* Foliis alternis.

65. LINUM USITATISSIMUM.

CHAR. SPEC. Calycibus capsulisque mucronatis, petalis crenatis, foliis lanceolatis alternis, caule subsolitario. **ANGLICE**, Flax.

LOCUS. Pannonia, Hispania. **Annua.**

VIS. Semina inodora, mucilagine, et oleo abundantia.

USUS. Semina in aqua infusa mucilaginis copiam præbent, et idcirco hoc modo usurpantur in Gonorrhœâ, ardore urinæ, et Catarrho: eorum pulvis cataplasma commune suppeditat.

COMP. Inf. Lini. *Lond.* Oleum Lini. *Lond. Edin. Eblan.*

**** Foliis opposits.**

66. LINUM CATHARTICUM.

CHAR. SPEC. Foliis oppositis ovato-lanceolatis, caule supernè dichotomo, corollis acutis.

ANGLICE, Purging Flax.

VIS. Amara, purgans.

CLASSIS VI.

ORDO I.

HEXANDRIA. MONOGYNIA.

***** Flores Spathacei sive Glumacei.**

ALLIUM. Willd. Spec. Plant. ii. 63. Persoon Syn. i. 355.

ORDO NAT. CLASSIS III. PLANTÆ MONOCOTYLEDONES STAMINA PERIGYNA.

ORDO VI. ASPHODELI.

V. Flores umbellati. Radix bulbosa. Calyx sex-partitus æqualis. **Juss.**

CHAR. GEN. Corolla sex-partita, patens. Spatha

multiflora. Umbella congesta. Capsula supera. (Flores capitati aut umbellati.)

* Folia caulina, plana. Umbella capsulifera.

67. ALLIUM PORRUM.

CHAR. SPEC. Caule planifolio umbellifero, staminibus tricuspidatis, radice tunicata, foliis basi vaginantibus.

ANGLICE, Leek.

LOCUS. Helvetia. Biennis.

VIS. Stimulans.

Oleo acri graveolente scatet, coctione fugatum. Pulpa, vel succus cutem stimulat, ad vesicationem usque.

** Folia caulina plana. Umbella bulbifera.

68. ALLIUM SATIVUM.

CHAR. SPEC. Caule planifolio bulbifero, bulbo composito, staminibus tricuspidatis.

ANGLICE, Garlic.

LOCUS. Sicilia. Perennis.

VIS. Stimulans.

**** Folia radicalia. Scapus nudus.

69. ALLIUM CEPA.

CHAR. SPEC. Scapo nudo inferne ventricosus, longiore, foliis teretibus.

ANGLICE, Onion.

**** *Flores nudi.*

ALOE. Willd. Spec. Plant. ii. 184. Persoon Syn. i. 378.

ORDO NAT. CLASSIS III. ORDO VI. ASPHODELI. I. Flores Spicati. Radix fibrosa. Calyx tubulosus. Juss.

CHAR. GEN. Corolla tubulosa, ore subsexfido patulo fundo nectarifero. Filamenta receptaculo inserta. Capsula supera, trilocularis trivalvis, polysperma. Semina biserialia margine membranaceo. (Floris ore nunc recto nunc labiato. Folia carnosa sive succulenta crassa.)

70. ALOE SPICATA.

CHAR. SPEC. Caulescens, foliis planis ensiformibus dentatis, floribus spicatis, companulatis horizontalibus.

ANGLICE, Aloes.

LOCUS, Caput Bonæ Spei, regionibus interioribus. Optimam Aloem habemus ex Insula Socotoræ prope Freta Babelmandel. Perennis.

71. ALOE VULGARIS, (flava.)

CHAR. SPEC. Foliis patulis lanceolatis margine spinosis, scapo ramoso, ramis bractea duplici amplexis.

LOCUS. India. Perennis.

VIS. Amarissima. Aloe Socotorina fusci-rubicundi coloris margine translucens; attamen inter se valde differt, fracturâ, colore, odore, sed parum effectum medicinali; sæpe Resinâ fraudulenter liquefacta, quæ aquâ bulliente, de more, intacta perstat. Viribus potentioribus hoc medicamen insigniter notatum, ut catharticum, emmenogogum, anthelminticum: super intestina inferiora actionem exercens, quæ plenè sed lentè evacuat, et amaritudine stomachum interdum renovat.

USUS. Perutilis Hypochondriasi, Dyspepsia, Chlorosi, et morbis Uteri inertia orientibus; siquidem Uterum æque ac Rectum stimulat, usurpari nunquam debet in Hæmorrhoida et Menorrhagiâ.

Aloe in aqua soluta cum Potassæ Subcarbonate enema efficax constituit ad ascarides arcendas.

DOSIS. Gr. ij. ad xv.

COMP. Pulv. Aloes Comp. *Lond.* Pil. Aloes. *Edin. Eblan.* Pil. Al. Comp. *Lond.* Pil. Aloes cum Myrrha. *Lond. Edin. Eblan.* Dec. Aloes Comp. *Edin. Eblan.* et alia plura.

SCILLA. Willd. Spec. Plant. ii. 125. Persoon Syn. i. 365.

ORDO NAT. CORONARIÆ. Linn.

ASPHODELI. Juss.

CHAR. GEN. Corolla hexapetala, patens, decidua. Filamenta filiformia, petalorum basi adnexa.

72. SCILLA MARITIMA.

CHAR. SPEC. Nudiflora, bracteis refractis. Folia lanceolata stricta. Scapus ante folia, longissimus multiflorus. Flores albi.

ANGLICE, Squill.

LOCUS. Hispaniæ, Lusitaniæ, Siciliæ, Syriæ littora arenosa, et in Barbariæ agris: bulbus crassitudinem fere capitis humani adæquat. Perennis.

VIS. Acris, amara, *cruda* corrosiva continet principium acre, album translucens, a D. Vogel detectum, Scillatin nuncupatum.

Dosibus amplis vomitum, et purgationem, minoribus autem expectorationem, et diuresin, ciet. Pneumoniâ aliisque morbis pyrexiâ comitatis, usus Scillæ parum convenit, propter acritudinem; quare Ipecacuanha longe anteposenda est. Hydrargyro conjuncta urinæ secretionem acriter auget, in Anasarcâ, Ascitide, et Hydrothorace.

DOSIS. Gr. i. ad v. ter die. Tinctura ʒss. Nimiâ quantitatē adhibitâ, vomitum, purgationem, stranguriam, dejectiones cruentas, gastrodyniam excitat.

COMP. Acetum Scillæ. *Lond. Edin. Eblan.* Oxy-mel Scillæ. *Lond. Eblan.* Pil. Scillæ Comp. *Lond. Edin. Eblan.* Pulv. Scillæ. *Edin. Eblan.* Syrupus Scillæ maritimæ. *Edin.* Tinct. Scillæ. *Lond. Eblan.*

***** Flores Incompleti.

ACORUS. Willd. Spec. Plant. ii. 199. Persoon Syn. i. 382.

ORDO NAT. PIPERITA. Linn.

CLASSIS II. PLANTÆ MONOCOTYLEDONES.
STAMINA HYPOGYNA. ORDO I. AROIDEÆ.
II. Spadix nudus, spathâ destitutus. Juss.

CHAR. GEN. Spadix cylindricus, tectus flosculis. Corolla hexapetala, nuda. Stylus nullus. Capsula trilocularis.

73. ACORUS CALAMUS.

CHAR. SPEC. Scapi mucrone longissimo foliaceo.

ANGLICE, Sweet-flag.

LOCUS. Europæ, et Indiæ fossis. Perennis.

VIS. Aromatica, acris, amaricans, fragrans.

CLASSIS VI.

ORDO III.

HEXANDRIA. TRIGYNIA.

* *Flores Inferi,*

COLCHICUM. Wild. Spec. Plant. ii. 272. Persoon
Syn. i. 398.

ORDO NAT. DENUDATÆ. Linn.

CLASSIS III. PLANTÆ MONOCOTYLEDONES.

STAMINA PERIGYNA. ORDO III. JUNC. IV.

Germina plura sæpius tria. Capsulæ totidem uniloculares,
interdum basi coalitæ. Flores paniculati aut spicati.
Juss.

CHAR. GEN. Spatha. Corolla sex partita: tubo ra-
dicato. Anthoræ incumbentes. Capsulæ tres, connexæ,
inflatæ.

74. COLCHICUM AUTUMNALE.

CHAR. SPEC. Foliis planis lanceolatis, erectis, petalis
ovatis. Flores autumnæ, et folia unâ cum fructu vere
profert.

ANGLICE, Meadow Saffron.

LOCUS. Prata humidiora. Perennis.

VIS. Acris, amara, diuretica, purgans. In Arthritide et
Rheumatismo colchicum quasi specificum habendum est;
paroxysmum amovens, nunc purgationem, nunc diuresin aut
diaphoresin excitans, et rarò sine claro, et aperto effectu.
Sed fatendum est, insultus podagricos crebriores fieri usu

medicamenti diurno. Quando versus intestina actionem suam dirigit, dejectiones copiosæ biliosæ sæpe producuntur.

DOSIS. Gr. iij. ad ix. pulv. siccati. Tinct. minima xxx. ad xl.

COMP. Acetum Colchici. *Lond.* Oxymel Colchici. *Eblan.* Syr. Colch. Autumnalis. *Edin.* Vinum Sem. Colch. Sp. Colchici. *Ammon. Lond.*

RUMEX. Willd. Spec. Plant. ii. 249. Persoon Syn. i. 394.

ORDO NAT. VAGINALES. Linn.

CLASSIS VI. PLANTÆ DICOTYLEDONES APETALÆ. STAMINA PERIGYNA. ORDO V. POLYGONEÆ. Juss.

CHAR. GEN. Calyx triphyllus. Petala tria conniventia. Semen unicum triquetrum.

** Valvulis granulo destitutis sive nudis.

75. RUMEX AQUATICUS.

CHAR. SPEC. Valvulis integerrimis, nudis, foliis latè cordatis glabris acutis.

ANGLICE, Great Water Dock.

LOCUS. Ad fluviorum, et paludum ripas. Perennis.

VIS. Sabstringens.

*** Floribus declinis.

76. RUMEX ACETOSA.

CHAR. SPEC. Floribus dioicis, caule elongato, foliis oblongis sagittatis.

ANGLICE, Sorrel.

LOCUS. Pascua, Sylvatica. Perennis.

VIS. Folia Potassæ bin-oxalatam continent quapropter gustum acidum habent.

CLASSIS VII.

ORDO I.

HEPTANDRIA. MONOGYNIA.

* *Flores completi.*

ÆSCULUS. Willd. Spec. Plant. i. 285. Persoon Syn. i. 403.

ORDO NAT. TRIHILATÆ. Linn.

CLASSIS XIII. PLANTÆ DICOTYLEDONES
POLYPETALÆ. STAMINA HYPOGYNA. ORDO
VI. ACERA. I. Fructus multilocularis. Juss.

CHAR. GEN. Calyx monophyllus, ventricosus. Corolla tetra-vel pentapetala inæqualis, pubescens, calyci inserta. Capsula trilocularis. Semina magna castaneæformia.

* Capsulis echinatis, corolla pentapetala planiuscula.

77. ÆSCULUS HIPPOCASTANUM.

CHAR. SPEC. Foliis digitatis septenis, corollis pentapetalis patulis.

ANGLICE, Horse Chesnut.

LOCUS. In Asia septentrionaliore. Perennis.

VIS. Cortex tonicus existimatur. Febribus intermittentibus.

CLASSIS VIII.

ORDO I.

OCTANDRIA. MONOGYNIA.

* *Flores completi.*

AMYRIS. Willd. Spec. Plant. ii. 333. Persoon Syn. i. 414.

ORDO NAT. CLASSIS XIV. PLANTÆ DICO-
TYLEDONES POLYPETALÆ. STAMINA PERI-
GYNA. ORDO XII. TEREBINACEÆ. II. Germen
simplex. Fructus multilocularis, loculis quibusdam interdum
abortivis. Juss.

CHAR. GEN. Calyx quadri-dentatus. Petala quatuor
oblonga patentia. Stigma subcapitatum. Bacca drupacea,
monosperma, per abortum.

78. AMYRIS ELEMIFERA.

CHAR. SPEC. Foliis ternatis quinato-pinnatisque sub-
tus tomentosis.

ANGLICE, Elemi-tree.

LOCUS. Carolina, Brazilia. Perennis.

VIS. Fragrans, aromatica. Resina Elemi dicta inest
Unguento Elemi Comp. *Lond. Eblan.*

79. AMYRIS GILEADENSIS.

CHAR. SPEC. Foliis ternatis integerrimis, pedunculis
unifloris lateralibus.

ANGLICE, Balsam of Gilead tree.

LOCUS. Arabia Felix. Perennis. Inusitata.

** Flores Incompleti.

DAPHNE. Willd. Spec. Plant. ii. 415. Persoon Syn.
i. 434.

ORDO NAT. VEPREULÆ. Linn.

CLASSIS VI. PLANTÆ DICOXYLEDONES APE-
TALE STAMINA PERIGYNA. ORDO II. THY-
MELÆÆ. Juss.

CHAR. GEN. Calyx nullus. Corolla quadrifida co-
rollacea, marcescens. Stamina includens. Drupa mono-
sperma.

* Floribus lateralibus.

80. DAPHNE MEZEREUM.

CHAR. SPEC. Floribus sessilibus, ternis caulinis, foliis
lanceolatis deciduis.

ANGLICE, Common Mezereon.

LOCUS. Anglia, et Europa septentrionalis. Perennis.

VIS. Acris, corrosiva.

USUS. Anti-rheumatica, anti-scorbutica, anti-syphilitica, existimatur.

COMP. Dec. Daphnes Mezerei. *Edin.* Dec. Sarsæ
Comp. *Lond.*

CLASSIS VIII.

ORDO III.

OCTANDRIA. TRIGYNIA.

POLYGONUM. Willd. Spec. Plant. ii. 440. Persoon Syn. i. 439.

ORDO NAT. VAGINALES. Linn.

CLASSIS VI. ORDO V. POLYGONEÆ. Juss.

CHAR. GEN. Calyx nullus. Corolla quinquepartita, calycina. Semen unicum, angulatum tectum. (Stamina et Styli numero incerti.)

*** Floribus spicatis, speciebus teretibus.

81. POLYGONUM BISTORTA. (Radice formâ.)

CHAR. SPEC. Caule simplicissime monostachyo, foliis ovatis in petiolum decurrentibus.

ANGLICE, Bistort.

LOCUS. In montibus, et pratis humidis. Perennis.

VIS. Styptica, astringens. Tannin, et acidum Gallicum continens.

CLASSIS IX.

ORDO I.

ENNEANDRIA. MONOGYNIA.

LAURUS. Willd. Spec. Plant. ii. 477. Persoon Syn. i. 448.

ORDO NAT. VAGINALES. Linn.

CLASSIS VI. ORDO IV. LAURI. Juss.

CHAR. GEN. Calyx, quadri-vel sexpartitus. Nectarium glandulis tribus, bisetis, germen cingentibus. Filamenta interiora quorum tria alterna sterilia glandulifera. Drupa monosperma. (Stamina 3, 5, 6, 7, 8, et 14. Flores passim dioici.)

82. LAURUS CINNAMOMUM.

CHAR. SPEC. Foliis suboppositis trinerviis ovato-oblongis, nervis versus apicem evanescentibus, panícula terminali.

ANGLICE, Cinnamon Tree.

LOCUS. Zeylona, Sumatra, Malabarìa, Martinica. Perennis.

VIS. Acris, dulcis, suavissima, aromatica, fragrantissima.

COMP. Aqua Cinnamomi. *Lond. Edin. et Eblan.*
Spiritus Cinnamomi. *Lond. Edin. et Eblan.* Tinct.
Cinnam. *Lond. Edin.* et *Eblan.* Tinct. Cinnam. Comp.
Lond. Pulv. Cinnam. Comp. *Lond. et Edin.*

83. LAURUS CASSIA.

CHAR. SPEC. Foliis triplinerviis lanceolatis.

ANGLICE, Cassia.

LOCUS. Idem ac supra. Perennis.

COMP. Aqua Lauri Cassiæ. *Edin.*

84. LAURUS CAMPHORA.

CHAR. SPEC. Foliis subtriplinerviis, lanceolato-ovatis, paniculis lateralibus.

ANGLICE, Camphor.

LOCUS. In Japonia. Perennis.

VIS. Aromatica, acris, amaricans, fragrantissima, stimulans. Narcotica? Sedativa?

Usurpari solet in Febribus continuis, et exanthematibus malignis, nec non in Epilepsiâ, Hysteriâ, et in plurimis morbis spasmodicis. Pro linimento adhibita est in Rheumatismo, et doloribus musculos afficientibus.

Camphoram non solum in Lauro Camphora, sed multis aliis vegta bilibus inesse compertum est.

DOSIS. Gr. ij. ad. ði. dosibus plenioribus vomitum, vertiginem, et alia molesta inducit.

COMP. Mist. Camphoræ. *Lond. et Eblan.* Emulsio Camphorata. *Edin.* Spiritus Camphoræ. *Lond. Edin. et Eblan.* Tinct. Camph. Comp. *Lond. Edin. et Eblan.* et Linimenta varia.

85. LAURUS NOBILIS.

CHAR. SPEC. Foliis lanceolatis coriaceis subundulatis venosis perennantibus, floribus subquadrifidis dioicis.

ANGLICE, Sweet-Bay.

LOCUS. Italia, Græcia. Perennis.

VIS. Calida, aromatica, fragrans.

86. LAURUS SASSAFRAS.

CHAR. SPEC. Foliis deciduis integris trilobisque, floribus dioicis. Stamina floris masculi novem, fæminei sex, brevissima.

ANGLICE, Sassafras.

LOCUS. Pennsylvania, Florida, Virginia. Perennis.

VIS. Acris, aromatica, fragrans.

COMP. Oleum Sassafras. *Lond. Edin. et Eblan.* Dec. Sarsæ. Comp. *Lond. Eblan.* Dec. Guaiaci. *Lond. Edin. et Eblan.* Aqua Calcis. Comp. *Eblan.*

CLASSIS IX.

ORDO III.

ENNEANDRIA. TRIGYNIA.

RHEUM. Willd. Spec. Plant. ii. 488. Persoon Syn.
i. 451.

ORDO NAT. VAGINALES. Linn.

CLASSIS VI. ORDO V. POLYGONEÆ. Juss.

CHAR. GEN. Calyx nullus. Corolla sexfida, persistens.
Semen unicum triquetrum.

87. RHEUM UNDULATUM.

CHAR. SPEC. Foliis subvillosis, petiolis æqualibus.

ANGLICE, Waved-leaved Rhubarb.

LOCUS. Chinæ ad murum, Siberia. Perennis.

88. RHEUM PALMATUM.

CHAR. SPEC. Foliis palmatis acuminatis.

ANGLICE, Rhubarb.

LOCUS. China. Perennis.

VIS. Nauseosa, styptica, amara, lutea. Purgans tonica
astringens.

Rhei radix ex sequentibus constat principiis, auctore
D. Brande.

Aquæ	- - - - -	8.	2.
Gummi	- - - - -	31.	0
Resinæ	- - - - -	10	0
Extracti, Tannin, Acidi gallici.	- -	26	0
Calcis Phosphatis	- - - - -	2	0
Calcis Malatis	- - - - -	6.	5.
Fibræ Lignæ	- - - - -	16.	3.

100 0

Alii Philosophi Chemici certiores nos faciunt; copiam
calcis Oxalatis huic radici inesse.

Alvo prius bene solutâ, hujus remedii ope preciosissimi, intestina postea leniter astringuntur, unde ejus in Dysenteria et Diarrhæâ summum beneficium. In febrium ultimo gradu malignarum, quando intestina evacuare oportet, Rhei radix ob predictam rationem, actionem idoneam inducit, citra periculum nimii alvi profluvii concitandi. In dyspepsiâ non solum alvum laxat, sed quoque stomachi vires debilitas insignitè roborat. Plurimis morbis Rheum cum aliis purgantibus feliciter conjungi potest.

Mirum sanè et observatione optimè dignum, quam citissimè urinam proprio colore tingit, quod sæpe accidit etiam intra sextam horæ partem.

DOSIS. Gr. i. ad ʒss.

COMP. Infusum Rhei. *Lond. et Edin.* Vinum Rhei *Edin.* Tinct. Rhei. *Lond. Edin. Eblan.* Tinct. Rhei. *Comp. Lond.* Extractum Rhei. *Lond.*

CLASSIS X.

ORDO I.

DECANDRIA. MONOGYNIA.

* *Flores polypetalis irregulares.*

MYROXYLON. Willd. Spec. Plant. ii. 546. Persoon Syn. i. 462.

ORDO NAT. PLANTÆ INCERTÆ SEDIS. APETALÆ DICLINES GERMINE SUPERO. Juss.

CHAR. GEN. Calyx campanulatus, quinque dentatus, petala quinque, supremo reliquis majore. Germen corollâ longius, legumen apice monospermum.

89. MYROXYLON PERUIFERUM.

CHAR. SPEC. Foliis abruptè pinnatis bijugis, foliolis suboppositis. Arbor pulcherrima. Racemi axillares. Calyx viridi-canescens, cinctus extra orificium petalis antherisque albis, intra continens legumen viride, quod singularem faciem representat.

ANGLICE, Sweet-smelling Balsam tree.

LOCUS. America meridionalis. Perennis.

VIS. Tenax, flavescens, amaro-acris, suaveolens.

Raro nisi extrinsecus usurpatur ulceribus mali moris.

Continet circiter decimam acidi Benzoici partem, cui præcipue tribuuntur vires, monente D. Stolze.

CASSIA. Willd. Spec. Plant. ii. 513. Persoon Syn. i. 456.

ORDO NAT. LOMENTACEÆ. Linn.

CLASSIS XIV. ORDO XI. LEGUMINOSÆ. II.

Corolla regularis. Legumen uniloculare bivalve. Stamina decem distincta. Arbores aut frutices; folia abruptè pinnata. Juss.

CHAR. GEN. Calyx pentaphyllus. Petala quinque, æqualia. Antheræ supremæ tres, steriles; infimæ tres rostratæ, in filamentis longioribus incurvis. Lomentum planum.

90. CASSIA SENNA.

CHAR. SPEC. Foliis sejugis subovatis, petiolis eglandulatis.

ANGLICE, Senna.

LOCUS. In Italiæ, et Indiæ satis, et præcipue in Ægypto superiore. Perennis.

VIS. Amara, nauseosa. Purgans.

Sæpenumero folia Cynanchi et Coronillæ fraudulentè commiscentur inter Sennæ sinceræ folia. DD. Lassaigne et Feneulle materiem Sennæ propriam, se nuper invenisse existimarunt, cui planta debet vim purgativam, ideoque ei imposuere nomen Cathartin, sed de hac re non satis constat. Senna per se adhibita tormina sæpe excitat, idcirco medicamentis carminativis adjungi debet, et alvus sic bene elicitur.

DOSIS. Pulveris Foliorum \mathfrak{z} i. ad \mathfrak{z} i. Infusi \mathfrak{z} ss. ad \mathfrak{z} ij.

COMP. Confectio Sennæ Comp. Lond. Edin. et Eblun. Ext. Cassiæ Sennâ. Edin. Inf. Sennæ. Lond. Eblan. Inf. Tamarindi cum Sennæ. Edin. et Eblan. Pulv. Sennæ

Comp. *Lond.* Tinct. *Sennæ. Lond. Edin. et Eblan.* Tinct. *Sennæ* Comp. *Edin.* Syrupus *Sennæ. Edin.*

CATHARTOCARPUS. Persoon Syn. i. 459.

CHAR. GEN. Calyx quinque partitus, deciduus. Corolla regularis, pentapetala. Filamenta inferiora arcuata. Legumen longum teres, lignosum, pluriloculare: loculis pulpâ fartia.

91. CATHARTOCARPUS FISTULA. Persoon.

CHAR. SPEC. Foliis quinque jugis ovatis acuminatis, petiolis eglandulosis. Petalis planis ovatis.

ANGLICE, Purging Cassia, or Cassia Fistula.

LOCUS. In India, Ægypto, et America. Perennis.

VIS. Dulcis, subnauseosa. Pulpa laxativa. Refrigerans, laxans. Sed ventris inflationem magnam inducit.

COMP. Pulpa Cassiæ Fistularis expressa. *Edin.* Conf. Cassiæ. *Lond. Edin. et Eblan.* Conf. *Sennæ. Lond. Edin. et Eblan.*

** Flores polypetali æquales.

HÆMATOXYLON. Willd. Spec. Plant. ii. 547. Persoon Syn. i. 462.

ORDO NAT. LOMENTACEÆ.

CLASSIS XIV. ORDO XI. LEGUMINOSÆ. II.

Corolla regularis. Legumen uniloculare bivalve. Stamina decem distincta. Arbores aut Frutices, folia abruptè pinnata. Juss.

CHAR. GEN. Calyx quinque-partitus. Petala quinque. Capsula lanceolata, unilocularis, trivalvis, valvis navicularibus.

92. HÆMATOXYLON CAMPECHIANUM.

ANGLICE, Logwood.

LOCUS. In Campechiæ Sinu. Perennis.

VIS. Styptica, tinctoria. Astringens in Dysenteria et Hæmorrhagiâ passivâ usurpata.

COMP. Ext. Hæmatoxyli. *Lond.*

SWIETENIA. Willd. Spec. Plant. ii. 557. Persoon Syn. i. 469.

CLASSIS XIII. ORDO XI. MELIÆ. III. Genera
Meliis affinia. Juss.

CHAR. GEN. Calyx quinquefidus. Petala quinque. Nectarium cylindricum, ore antheras gerens. Capsula tri-vel quinque-locularis, lignosa, magna, tri-vel quinque-valvis. Semina imbricata alata.

93. SWIETENIA FEBRIFUGA.

CHAR. SPEC. Foliis pinnatis subquadrijugis, foliolis elliptico-subrotundis emarginatis, basi inæqualibus, panicula terminali divaricata.

ANGLICE, Febrifuge Swietenia.

LOCUS. In montibus Indiæ Orientalis. Perennis.

VIS. Amara, astringens, tonica.

GUAIAACUM. Willd. Spec. Plant. ii. 538. Persoon Syn. i. 462.

CLASSIS XIII. ORDO XXI. RUTACEÆ. I. Folia
stipulacea sæpius opposita. Juss.

CHAR. GEN. Calyx quinquefidus inæqualis. Petala quinque, calyci inserta. Capsula angulata tri-sive quinque-locularis.

94. GUAIAACUM OFFICINALE.

CHAR. SPEC. Foliolis bijugis obtusis. Capsulæ bi-angulares.

ANGLICE, Guaiacum.

LOCUS. Hispania, Jamaica, Antillæ. Perennis.

VIS. Aromatica, subamara. Contra Rheumatismum chronicum usurpatum.

DOSIS. Gr. x. ad ʒss.

COMP. Dec. Guaiaci Comp. *Edin.* Mist. Guaiaci. *Lond.* Tinct. Guaiaci. *Lond. Edin.* et *Eblan.* Dec. Sarsæ Comp. *Lond.* et *Eblan.* Tinct. Guaiaci Ammon. *Lond. Edin.* et *Eblan.* Pulv. Aloes Comp. *Lond.* et *Eblan.*

RUTA. Willd. Spec. Plant. ii. 542. Persoon Syn. i. 464.

CLASSIS XIII. ORDO XXI. RUTACEÆ. II. Folia alterna nuda. Juss.

CHAR. GEN. Calyx quadri-sive quinque-partitus. Petala concava. Receptaculum punctis melliferis decem cinctum. Capsula lobata. (Petala quibusdem quatuor, et stamina octo.)

95. RUTA GRAVEOLENS.

CHAR. SPEC. Foliis suprà decompositis, foliolis oblongis, terminali obovato, petalis integerrimis. Flos primarius quinquefidus, reliqui quadrifidi.

ANGLICE, Rue.

LOCUS. Europa Australior, Alexandria, Mauritania. Perennis.

VIS. Stimulans, antispasmodica.

QUASSIA. Willd. Spec. Plant. ii. 567. Persoon Syn. i. 465.

CLASSIS XIII. ORDO XV. MAGNOLIÆ. Genera Magnoliis affinia. Juss.

CHAR. GEN. Calyx pentaphyllus. Petala quinque. Nectarium pentaphyllum. Drupæ quinque, distantes, bivalves, monospermæ, receptaculo carnosio insertæ. (Flores etiam monoici aut pentandri.)

96. QUASSIA SIMAROUBA.

CHAR. SPEC. Floribus paniculatis monóicis, foliis abruptè pinnatis, foliolis alternis subpetiolatis, petiolo nudo. Paniculæ axillares amplæ. Drupæ ovatæ, subnigræ, receptaculo carnosio lateralitèr affixæ.

ANGLICE, Simarouba, or Mountain Damson.

LOCUS. Locis arenosis Americæ australioris. Perennis.

VIS. Amara, tonica.

USUS. In Diarrhœâ, Dysenterîâ, Dyspepsiâ.

DOSIS. ʒi. ad ʒss.

COMP. Infusum Simaroubæ. Lond.

97. QUASSIA EXCELSA.

CHAR. SPEC. Floribus polygamis pentandris paniculatis, foliis impari-pinnatis, foliolis oppositis petiolatis, petiolo nudo.

ANGLICE, Quassia.

LOCUS. In submontosis Jamaicæ, et Caribæarum. Perennis.

VIS. Amarissima, inodora, subflava.

USUS. Quemadmodum Quassiæ Infusum, nec sales metallicos præcipitat, nec cum ferreis nigrescit, idoneum constituit vehiculum pro istis medicamentis adhibendis.

Infusum fortius saccharo commistum, optimum suppeditat venenum ad muscas interimendas.

COMP. Infusum Quassiæ. *Lond.* Tinct. Quassiæ Excelsæ. *Edin.* et *Eblan.*

*** Flores monopetali æquales.

RHODODENDRON. Willd. Spec. Plant. ii. 603. Persoon Syn. i. 478.

CLASSIS IX. PLANTÆ DICOTYLEDONES MONOPETALÆ. COROLLA PERIGYNA. ORDO II. RHODODENDRA. I. Corolla monopetala. Juss.

CHAR. GEN. Calyx quinque partitus. Corolla subinfundibuliformis, subobliqua. Stamina declinata. Capsula quinquelocularis.

98. RHODODENDRON CHRYSANTHUM.

CHAR. SPEC. Foliis oblongis scabris, subtus discoloribus glabris, umbellis terminalibus, corollis rotatis, petalis obovatis, irregularibus. Frutex parvus diffusus. Folia venoso-reticulata. Rami squamis stipuliformibus ferrugineis tecti. Flos luteus.

ANGLICE, Golden-flowered Rhododendrum.

LOCUS. Siberiæ alpebus. Perennis.

VIS. Astringens, subamara, stimulans, diaphoretica?

USUS. Rheumatismo, Arthritide.

ARBUTUS. Willd. Spec. Plant. ii. 616. Persoon Syn. i. 482.

ORDO NAT. BICORNES. Linn.

CLASSIS IX. ORDO III. ERICÆ. I. Germen superum. Juss.

CHAR. GEN. Calyx quinque partitus. Corolla ovata basi pellucida. Bacca supera, quinque locularis.

99. ARBUTUS UVA URSI.

CHAR. SPEC. Caulibus procumbentibus, foliis integerrimis. Baccæ pentaspermæ.

ANGLICE, Trailing Arbutus, or Bear-berry.

LOCUS. Sueciæ colles sabulosi, Canada. Perennis.

VIS. Astringens, dulcamara.

USUS. Morbis calculosis et Renalibus; Catarrho vesicæ.

DOSIS. Pulveris foliorum ʒi. ad ʒi.

STYRAX. Willd. Spec. Plant. ii. 623. Persoon Syn. i. 483.

ORDO NAT. HESPERIDÆ. Linn.

CLASSIS IX. ORDO I. GUAIACANÆ. I. Stamina definita. Juss.

CHAR. GEN. Calyx campanulatus subinteger, inferus. Corolla quinque vel septempartita: tubo brevi calyci inserta. Drupa mono-vel disperma, Coriacea. (Stamina, 6—16, corollæ tubo inserta. Antheræ oblongæ lineares.)

100. STYRAX OFFICINALE.

CHAR. SPEC. Foliis ovatis subtus villosis, racemis simplicibus, folio brevioribus.

ANGLICE, Storax.

LOCUS. Syria, Judæa, Italia. Perennis.

VIS. Aromatica, fragrans, stimulans, haud per se adhibita.

COMP. Styra^x Purificata. *Eblan*. Pil. e Styra^ce. *Eblan*.

101. STYRAX BENZOIN.

CHAR. SPEC. Foliis oblongis acuminatis subtus tomentosis, racemis compositis longitudine foliorum.

ANGLICE, Benjamin tree.

LOCUS. Sumatra, insula. Perennis.

VIS. Fragrans, pungens. Acidum Benzoicum adipiscitur Resinæ sublimatione.

COMP. Tinct. Benzoës Comp. *Lond. Edin. et Eblan.*

**** Flores apetalæ sive incompleti.

COPAIFERA. Willd. Spec. Plant. ii. 630. Persoon Syn. i, 462.

CLASSIS XIV. ORDO XI. LEGUMINOSÆ. X. Corolla irregularis (interdum nulla.) Stamina decem distincta. Legumen capsulare uniloculare submonospermum, sæpius non dehiscent. Arbores aut frutices; folia impari pinnata aut simplicia; stipulæ a petiolo distinctæ, mox deciduæ. Juss.

CHAR. GEN. Calyx nullus. Petala quatuor. Legumen ovatum. Semen unicum, arillo ovato.

102. COPAIFERA OFFICINALIS.

ANGLICE, Copaiba.

LOCUS. In Brasiliâ, Antillis. Perennis.

VIS. Pungens, amara, subflavus, stimulan, diuretica. Copaibæ balsamum Alkohole, et Æthere solubile.

USUS. Omnium medicamentorum, quæ in usu communi sunt contra Gonorrhæam, Copaibæ balsamum inter optima ferè habendum; sed adhiberi oportet, dosibus ita amplis, ut alvum benè laxet; alioqui parum prodest. Datur etiam Leucorrhœâ laborantibus, sed haud æquo beneficio.

DOSIS. M. x. ad ʒss. bis vel ter quotidie.

CLASSIS X.

ORDO II.

DECANDRIA.. DIGYNIA

DIANTHUS. Willd. Spec. Plant. ii. 671. Persoon Syn. i. 493.

ORDO NAT. CARYOPHYLLEÆ. Linn.

CLASSIS XIII. ORDO XXII. CARYOPHYLLEÆ.

V. Calyx tubulosus. Stamina decem (alterna hypogyna, alterna sæpius epipetala.) Styli duo, aut tres, aut quinque. Juss.

CHAR. GEN. Calyx cylindricus, longus, coriaceus, monophyllus; basi squamis quatuor vel octo. Petala quinque unguiculata. Capsula cylindrica, unilocularis, apice dehiscens.

** Flores dispersi in caule ramoso.

103. DIANTHUS CARYOPHYLLUS.

CHAR. SPEC. Floribus solitariis, squamis calycinis subrhombeis brevissimis, petalis crenatis imberbibus, foliis lineari-subulatis canaliculatis glaucis.

ANGLICE, Clove-pink, or Gilly-flower.

LOCUS. Italiæ alpes. Perennis.

VIS. Suaveolens, fragrans, roseo-tinctoria.

COMP. Syrupus Dianthi Caryophylli. *Edin. Eblan.*

CLASSIS X.

ORDO V.

DECANDRIA. PENTAGYNIA.

OXALIS. Willd. Spec. Plant. ii. 772. Persoon Syn. i. 514.

ORDO NAT. SUCCULENTÆ. Linn.

CLASSIS XIII. ORDO XIII. GERANIA. Juss.

CHAR. GEN. Calyx pentaphyllus. Petala unguibus connexa. Stamina inæqualia, quinque breviora, exteriora basi connata. Capsula angulis elasticè dehiscens, pentagona. Semina subarillata.

*** Foliis ternatis.

† Acaulis scapo unifloro.

104. OXALIS ACETOSELLA.

CHAR. SPEC. Foliis ternatis obcordatis, radice dentata.

ANGLICE, Wood Sorrel.

LOCUS. Europæ sylvæ. Perennis.

VIS. Folia acidula sunt, quia continent Potassæ Binoxolatem, vulgo *Essential Salt of Lemons*.

CLASSIS XI.

ORDO I.

DODECANDRIA. MONOGYNIA.

ASARUM. Willd. Spec. Plant. ii. 838. Persoon Syn. ii. 1.

ORDO NAT. SARMENTACEÆ. Linn.

CLASSIS V. PLANTÆ DICOTYLEDONES APETALÆ. STAMINA EPIGYNA. ORDO I. ARISTOLOCHIÆ. Juss.

CHAR. GEN. Calyx quadrifidus, germi insidens. Corolla nulla. Capsula coriacea, coronata.

105. ASARUM EUROPÆUM.

CHAR. SPEC. Foliis reniformibus obtusis binis.

ANGLICE, Asarabacca.

LOCUS. Europa Australior. Scanie in nemoribus, præsertim calcariis. Perennis.

VIS. Amaro-acris, nauseosa. Errhina.

COMP. Pulv. Asari Comp.

CANELLA. Willd. Spec. Plant. ii. 851. Persoon Syn. ii. 4

CLASSIS XIII. ORDO XI. MELIÆ. I. Folia simplicia. Juss.

CHAR. GEN. Calyx trilobus. Petala quinque. Antheræ sedecim, adnatæ nectario urceolato. Bacca unilocularis, di-sive tetra-sperma.

106. CANELLA ALBA.

ANGLICE, White, or laurel-leaved Canella.

LOCUS. In interioribus Americæ sylvis. Perennis.

VIS. Acris, pungens, subamara, aromatica, fragrans. Nunquam per se usitata.

DOSIS. Gr. x. ad ʒss.

COMP. Tinct. Gent. Comp. *Edin.* Vinum Aloes. *Lond. Edin. et Eblan.* Pulv. Aloes cum Canella. *Eblan.*

LYTHRUM. Willd. Spec. Plant. ii. 865. Persoon Syn. ii. 7.

ORDO NAT. CALYCANTHEMÆ. Linn.

CLASSIS XIV. ORDO IX. SALICARIÆ. I. Flores polypetalis. Juss.

CHAR. GEN. Calyx 6, 8, 10, 12 dentatus. Petala quinque vel sex, calyci inserta. Capsula tecta, bilocularis, polysperma. (Stamina, 2, 8, 9, 10, 12.)

107. LYTHRUM SALICARIA.

CHAR. SPEC. Foliis oppositis cordato-lanceolatis, floribus spicatis dodecandris. Flores verticillato-spicati.

ANGLICE, Loosestrife, or Purple willow herb.

LOCUS. Europæ aquosis indigena. Perennis.

VIS. Substringens.

USUS. In Diarrhæâ, et Dysenteriâ.

DOSIS. ʒss ad ʒiv.

CLASSIS XI.

ORDO II.

DODECANDRIA. DIGYNIA.

AGRIMONIA. Willd. Spec. Plant. ii. 875. Persoon Syn. ii. 9.

ORDO NAT. SENTICOSÆ. Linn.

CLASSIS XIV. ORDO X. ROSACEÆ. III. Germina plura definita (rariùs unicum,) calyce urceolari suprà coarctato tecta, quasi infera, singula monostyla. Semina totidem. Herbæ pleræque, plures apetalæ, plures definitè staminiferæ, quædam diclines. Juss.

CHAR GEN. Calyx quinque dentatus, altero obvallatus. Petala quinque. Semina duo, in fundo calycis.

108. **AGRIMONIA EUPATORIA.**

CHAR. SPEC. Foliis caulinis interruptè pinnatis: impari petiolato, fructibus hispidis.

ANGLICE, Common Agrimony.

LOCUS. Europæ prata. Perennis.

VIS. Styptica, diuretica.

USUS. Cachexia, Febres.

CLASSIS XII.

ORDO I.

ICOSANDRIA. MONOGYNIA.

CARYOPHYLLUS. Persoon Syn. ii. 30.

EUGENIA. Willd. Spec. Plant. ii. 959.

ORDO NAT. ARBUSTIVA. Linn.

CLASSIS XIV. ORDO MYRTI. I. Flores in foliorum axillis, aut in pedunculis multifloris oppositi. Folia plerumque opposita, et punctata. Juss.

CHAR. GEN. Calyx infundibuliformis, quadrifidus. Petala quatuor. Drupa sive bacca sicca, ovata, uni-vel bilocularis, laciniis calycis coronata.

109. CAROPHYLLUS AROMATICUS. Persoon.
EUGENIA CARYOPHYLLATA. Willd. Spec. Plant.

ii. 965.

ANGLICE, Clove.

LOCUS. In Moluccis, solo aridissimo. Perennis.

VIS. Aromatica, fragrans, pungens.

COMP. Inf. Caryophyllorum. *Lond.* et alia plura.

MYRTUS. Willd. Spec. Plant. ii. 967. Persoon Syn.

ii. 30.

ORDO NAT. ARBUSTIVA. Linn.

CLASSIS XIV. ORDO VII. MYRTI. Juss.

CHAR. GEN. Calyx quinquefidus, superus. Petala quinque. Bacca bi-vel tri-locularis polysperma.

110. MYRTUS PIMENTA.

CHAR. SPEC. Foliis alternis, oblongis.

ANGLICE, Allspice.

LOCUS. Jamaica. Perennis.

VIS. Aromatica, fragrans.

COMP. Aqua Pimentæ. *Lond. Edin. et Eblan.*

Ol. Piment. *Lond. Edin. et Eblan.* **Pil. Opiata.** *Edin.*

Syr. Rhamni. *Lond.*

PUNICA. Willd. Spec. Plant. ii. 981. Persoon Syn.

ii. 33.

ORDO NAT. POMACEÆ. Linn.

CLASSIS XIV. ORDO VII. MYRTI. Juss.

CHAR. GEN. Calyx quinquefidus, superus. Petala quinque. Bacca (Pomum. Linn.) multilocularis, polysperma. Receptacula parietalia. Semina baccata.

111. PUNICA GRANATUM.

CHAR. SPEC. Foliis lanceolatis, caule arboreo.

ANGLICE, Pomegranate.

LOCUS. Africa, Hispania, Italia. Perennis.

VIS. Styptica, rubra.

USUS. Diarrhæâ, Dysenteriâ. Decoctum quasi injectio in Leucorrhæâ, et Gargarisma in Cynanche.

DOSIS. Pulv. ʒss. ad ʒi. Decocti ʒss.

AMYGDALUS. Willd. Spec. Plant. ii. 982. Perseon Syn. ii. 33.

ORDO NAT. DRUPACEÆ. Linn.

CLASSIS XIV. ORDO X. ROSACEÆ. VII. Ger-
men unicum superum monostylum. Nux mono—aut dis-
perma, nuda aut sæpius Drupacea. Arborea aut frutices.
Amygdalæ. Juss.

CHAR. GEN. Calyx quinquefidus, inferus. Petala quinque. Drupa nuce poris perforata. Flores sessiles.

**** Drupa villosa, coriacea nuce porosa.**

112. AMYGDALUS COMMUNIS.

CHAR. SPEC. Foliorum serraturis infimis glandulosis, floribus sessilibus geminis.

ANGLICE, Common Almond.

LOCUS. Tripolis sepes. Perennis.

VIS. Dulcis, sapida.

β. AMYGDALUS AMARA.

ANGLICE, Bitter Almond.

VIS. Amara.

Amygdalorum ambæ varietates supra descriptæ, scilicet dulcium et amararum, ejusdem olei prope dimidium ponderis ipsarum *expressione* suppeditant; quod vulgo Oleum Amygdalæ dicitur. Porro varietas amara *distillatione* insuper præbet oleum essenziale, animalibus prorsus lethale; nam in eo continetur, auctore D. Hennell, acidum Hydrocyanicum, omnium fere rerum, hactenus cognitarum, citissimam mortem inducens.

COMP. Mist. Amygdalarum. Lond. Edin. et Eblan.

Emulsio Arabica. *Edin.* et *Eblan.* Emulsio Camphorata.
Edin. Confectio Amygdalarum. *Lond.*

PRUNUS. Willd. Spec. Plant. ii. 984. Persoon Syn.
 ii. 34.

ORDO NAT. DRUPACEÆ. Linn.

CLASSIS XIV. ORDO X. ROSACEÆ. Juss.

CHAR. GEN. Calyx quinquefidus, inferus. Petala
 quinque. Drupæ lævis nux suturis prominulis.

** Prunus. Drupa major, carnosa, nucleo subcompresso,
 ovato.

113. PRUNUS DOMESTICA.

-CHAR. SPEC. Pedunculis subsolitariis, foliis lance-
 olato-ovatis convolutis, ramis muticis.

ANGLICE, Plum.

LOCUS. Europa Australis. Perennis.

VIS. Acidula, laxativa.

COMP. Confectio Sennæ. *Lond. Edin.* et *Eblan.*

CLASSIS XII.

ORDO V.

ICOSANDRIA. PENTAGYNIA.

PYRUS. Willd. Spec. Plant. ii. 1012.

ORDO NAT. POMACEÆ. Linn.

CLASSIS XIV. ORDO X. ROSACEÆ. I. Germen
 simplex inferum, polystylum. Pomum calycino limbo umbi-
 licatum, multiloculare. Arbores aut frutices Pomaceæ. Juss.

CHAR. GEN. Calyx quinquefidus. Petala quinque.
 Pomum inferum, quinque-loculare, polyspermum.

114. PYRUS CYDONIA.

CYDONIA CYDONIA. Persoon Syn. ii. 40.

CHAR. SPEC. Foliis ovatis integerrimis, floribus solitariis.

ANGLICE, Quince.

LOCUS. Ad ripas petrosas Danubii. Perennis.

VIS. Acido-styptica. Semina mucilaginosa.

COMP. Decoctum Cydoniæ. *Lond.*

CLASSIS XII.

ORDO VIII.

ICOSANDRIA. POLYGYNIA.

ROSA. Willd. Spec. Plant. ii. 1063. Persoon Syn. ii. 47.

ORDO NAT. SENTICOSÆ. *Linn.*

CLASSIS XIV. ORDO X. ROSACEÆ. II. Germina plura indefinita, calyce urceolari suprà coarctato tecta, quasi infera, singula monostyla. Semina totidem. Frutices. *Rosæ. Juss.*

CHAR. GEN. Calyx urceolatus, quinquefidus, carnosus, collo coarctatus. Petala quinque. Semina plurima, hispida, calycis interiori lateri affixa.

** Germinibus ovatis.

115. ROSA CENTIFOLIA.

CHAR. SPEC. Pedunculis hispidis, caule hispido aculeato, petiolis inermibus, foliis subtus pubescentibus.

ANGLICE, Hundred-leaved Rose.

LOCUS. Europa, ubi ceteræ sequentes nascuntur. Perennis.

VIS. Fragrans, rubra.

COMP. Aqua Rosæ. *Lond. Edin. et Eblan. Syr. Rosæ. Lond. Edin.*

116. ROSA GALLICA.

CHAR. SPEC. Pedunculis hispidis, caule, petiolisque hispido-aculeatis.

ANGLICE, Red Rose.

VIS. Astringens, tonica.

COMP. Inf. Rosæ. *Lond. Edin. et Eblan.* Mel Rosæ. *Lond. et Edin.* Syr. Rosæ. *Edin.*

117. ROSA CANINA.

CHAR. SPEC. Pedunculis glabris, caule, petiolisque aculeatis. Foliis ovatis glabris. Petioli subvillosi.

ANGLICE, Dog Rose.

LOCUS. In sylvis, et ad sepes. Perennis.

VIS Fragrans, acidula.

COMP. Conf. Rosæ Caninæ. *Lond.*

TORMENTILLA. Willd. Spec. Plant. ii. 1112. Persoon Syn. ii. 56.

ORDO NAT. SENTICOSÆ. Linn.

CLASSIS XIV. ORDO X. ROSACEÆ. IV. Germina plura indefinita, verè supera, receptaculo communi imposita, singula monostyla. Semina totidem nuda, aut rariùs baccata. Herbæ, aut rariùs frutices. Potentillæ. Juss.

CHAR. GEN. Calyx 8-fidus. Petala quatuor. Semina subrotunda, nuda, receptaculo parvo, exsucco affixa.

118. TORMENTILLA ERECTA, VEL OFFICINALIS.

CHAR. SPEC. Caule erectiusculo, foliis sessilibus; foliolis ovato-lanceolatis serratis.

ANGLICE, Tormetil, or Septfoil.

LOCUS. Europæ pascua, et sylvæ. Perennis.

VIS. Styptica, rubra. Radix copiam Tannin continet.

USUS. In Diarrhœa quasi astringens.

DOSIS. Gr. x. ad ʒi. pulveris. Dec. ʒi.

COMP. Pulvis Cretæ Comp. *Lond.*

GEUM. Willd. Spec. Plant. ii. 1113. Persoon Syn. ii. 56.

ORDO NAT. SENTICOSÆ. Linn.

CLASSIS XIV. ORDO X. ROSACEÆ. Sectio IV.

Juss.

CHAR. GEN. Calyx 10-fidus, laciniae inæquales. Petala quinque. Semina barbata sive aristata; arista plerumque geniculata.

119. GEUM URBANUM.

CHAR. SPEC. Floribus erectis, aristis uncinatis nudis, foliis caulinis ternatis, radicalibus lyrato-pinnatis.

ANGLICE, Avens, or Herb Bennet.

LOCUS. Europæ nemoribus indigena. Perennis.

VIS. Fragrans. Caryophyllum redolens. Subamara.

USUS. Febribus intermittentibus. Dysenteria, Diarrhoea.

DOSIS. ʒss. ad ʒi. Pulv. Dec. ʒi.

CLASSIS XIII.

ORDO I.

POLYANDRIA. MONOGYNIA.

* *Tetrapetali.*

PAPAVER. Willd. Spec. Plant. ii. 1144. Persoon Syn. ii. 62.

ORDO NAT. RHÆADES. Linn.

CLASSIS XIII. ORDO II. PAPAVERACEÆ. I. Stamina indefinita; antheræ filamentis adnatæ. Juss.

CHAR. GEN. Calyx di-phyllus. Corolla tetra-petala. Capsula unilocularis, sub stigmate persistente poris dehiscens.

** Capsulis glabris.

120. PAPAVER RHÆAS.

CHAR. SPEC. Capsulis subglobosis, caule multifloro piloso; pilis patentibus, foliis pinnatifidis incis.

ANGLICE, Corn, or Red Poppy.

LOCUS. Europæ arvis indigena. Annu.

VIS. Amariuscula, rubeo-tinctoria.

USUS. Petala solum usui sunt pro Syrupo Rhœados conficiendo.

121. PAPAVER SOMNIFERUM.

CHAR. SPEC. Calycibus, capsulisque glabris, foliis amplexicaulibus incis. glaucis.

ANGLICE, White Poppy.

LOCUS. Asia, Europæ australioris agri. Annu.

VIS. Resinose-gummosa, amara, acris, tetra-foetida.

Galli meridionales magnam hujus plantæ copiam propter semina colunt, quorum expressione oleum fixum, blandum largitèr obtinent. Turcis mos est immaturas Papaverum capsulas longitudinalitèr secare, et succum lacteum vulneribus extemplò defluentem, solis radiis exponere, donec idoneam crassitudinem habeat, et postremò, sub placentarum formâ, nobis deportare. Eandem rationem persequuntur Indi Orientales, opium proferentes, Turcico longè præstantius. Sed in Anglia etiam, eodem modo, opium optimi generis acquisitum fuit, a DD. Ball, Cowley, Staines, et Young.

Hujus Medicamenti vires, proprio principio, naturâ alkalinâ prædito, attribuuntur, quod D. Sertürner auctor, morphiam vocavit. Hæc opio invenitur acido Meconico nuncupato consociata, quod per se nullam potestatem narcoticam exercet.

Apud ægrotos, hoc princeps anodynorum duplici ratione valere videtur, nempe stimulante aut sedante.

Dosis parva vim stimulantem clarè exercet, actionem systématis arterialis auget, mentem subindè solatur, vertiginem et pyrexiam inducit, quorum omnia quies, et somnus sequuntur. Post dosin ampliorem vis stimulans ægrè percipitur.

Opium omnes præter cutis secretionem imminuit, alvum astringit, et urinam parciorem efficit. Sic remedium posidemus sæpenumero perutile, ad dolorem sublevan-

dum, somnum inducendum, secretiones intemperatas cohibendas : et tam frequenter medicinæ praxi adhibitum, ut difficile esset omnes morbos indicare, quibus conveniat. Attamen frequentior est ejus usus in Febribus, Dysenteria, Diarrhœa, Cholera, Rheumatismo, Colica Pictonum, Spasmis; doloribus accerbissimis, nunquam verò ex inflammatione orientibus.

Observatione dignissimum est, opium, pro Toxico sump-tum, pupillæ oculi fines semper contrahere; quâ potestate ab Belladonna, et Hyoscyamo maximè differt.

DOSIS. Pulv. Opii gr. ss. ad gr. ij. Tinct. Opii. m. v. ad 3ss.

COMP. Conf. Opii. *Lond. et Edin.* Elect. Catechu. *Edin.* Ext. Opii. *Lond. Edin. et Eblan.* Pil. Opii. *Edin.* Pil. Saponis cum Opio. *Lond.* Pulv. Opiatus. *Edin.* Pulv. Ipecac. Comp. *Lond. Edin. et Eblan.* Tinct. Opii. *Lond. Edin. et Eblan.* et alia plura.

CLASSIS XIII.

ORDO III.

POLYANDRIA. TRIGYNIA.

DELPHINIUM, Willd. Spec. Plant. ii. 1226. Per-
soon Syn. ii. 82.

ORDO NAT. MULTISILIQUÆ, Linn.

CLASSIS XIII. ORDO I. RANUNCULACEÆ. II.
Capsulæ polyspermæ intus dehiscentes. Petala irregularia.
Calyx sæpè coloratus, a Linnæo corolla dictus, petalis ab
eodem in nectaria conversis. Juss.

CHAR. GEN. Calyx nullus. Petala quinque. Nec-
tarium bifidum, postice cornutum. Siliquæ tres sive una.

** Tricapsularia.

122. DELPHINIUM STAPHISAGRIA.

CHAR. SPEC. Nectariis tetraphyllis petalo brevioribus, foliis palmatis, lobis obtusis.

ANGLICE, Stavesacre.

LOCUS. Istria, Dalmatia, Calabria, Apulia. Biennis.

VIS. Nauseosa, acris, amaricans, emetica, cathartica.

DD. Lassaigue et Feneulle in *Staphisagriae* seminibus novum alkali invenerunt, cui nomen dederunt *Delphinium*, ex qua, plantae vires pendent.

USUS, Pediculos necare, inter stragula inspersa.

ACONITUM. Willd. Spec. Plant. ii. 1232. Persoon Syn. ii. 83.

ORDO NAT. MULTISILIQUEÆ. Linn.

CLASSIS XIII. ORDO I. RANUNCULACEÆ.

Sectio II. Juss.

CHAR. GEN. Calyx nullus. Petala quinque, supremo fornicato. Nectaria (cucullus) duo pedunculata, recurva. Siliquae tres sive quinque.

**** Corollis cæruleis.**

123. ACONITUM NAPELLUS.

CHAR. SPEC. Cuculli calcare recto obtuso, labio lanceolato ascendente bifido, galea convexa, foliis nitidis quinque partitis, laciniis tripartitis incisis linearibus.

ANGLICE, Monkshood, or Wolfsbane.

LOCUS. Europæ humidiuscula. Perennis.

VIS. Amara, acris, nauseosa, toxica. Narcotica, sudorifica existimata.

USUS. Rheumatismo.

DOSIS. Extracti gr. j. bis terve die.

ACONITUM NEOMONTANUM.

CHAR. SPEC. Cuculli calcare adunco obtuso, labio lanceolato revoluta bifido, galea convexa, foliis nitidis digitato-quinquepartitis, laciniis lato-lanceolatis brevissime dentatis, pedunculis glabris. Flores violacei, galea humili fornicata, breviter rostrata.

CLASSIS XIII.

ORDO IV.

POLYANDRIA. TETRAGYNIA.

WINTERA. Willd. Spec. Plant. ii. 1239. Persoon Syn. ii. 84.

CLASSIS XIII. ORDO XI. MELIÆ. I. Folia simplicia. Juss.

CHAR. GEN. Calyx trilobus. Petala sex, sive duodecim. Germina clavata. Styli nulli. Baccæ quatuor sive octo, obovatæ.

124. WINTERA AROMATICA.

CHAR. SPEC. Pedunculis axillaribus aggregatis subtrifloris, floribus tetragynis.

ANGLICE, Winter's Bark.

LOCUS. In Freti Megallanici vallibus. Perennis.

VIS. Acris, aromatica, tonica.

USUS. In Dyspepsiâ, Scorbuto.

CLASSIS XIII.

ORDO VI.

POLYANDRIA. POLYGYNIA.

HELLEBORUS. Willd. Spec. Plant. ii. 1335. Persoon Syn. ii. 107.

ORDO NAT. MULTISILIQUEÆ. Linn.

CLASSIS XIII. ORDO I. RANUNCULACEÆ. Juss.

CHAR. GEN. Calyx nullus. Petala quinque sive plura. Nectaria bilabiata, tubulata. Capsulæ quinque vel sex polyspermæ, erectiusculæ, compressæ.

125. HELLEBORUS NIGER.

CHAR. SPEC. Scapo subbifloro subnudo, foliis pedatis, foliolis cuneato-oblongis. Floret primo hyeme. Radix nigra. Flores albid.

ANGLICE, Black Hellebore.

LOCUS. In Pannoniæ, Hetruriæ, Apennini asperis. Perennis.

VIS. Acris, amaricans, nauseosa foetida. Cathartica Hydrogoga? Emmenogoga?

USUS. Olim in Mania, Melancholia, Hydrope, et mensium suppressione usurpata.

DOSIS. Gr. ij. ad x.

COMP. Tinct. Hellebori nigri. *Lond. Edin. Eblan.*
Ext. Hellebori nigri. *Edin. Eblan.*

126. HELLEBORUS FOETIDUS.

CHAR. SPEC. Caule multifloro folioso, foliis pedatis remote serratis coriaceis, corollâ subconnivente.

ANGLICE, Stinking Hellebore, or Bearsfoot.

LOCUS. In sylvis. Perennis.

VIS. Acris, nauseosa. Cathartica. Anthelmintica?

CLASSIS XIV.**ORDO I.****DIDYNAMIA. GYMNOSPERMIA.**

* *Calyces subquinquefidi.*

HYSSOPUS. Willd. Spec. Plant. iii. 47. Persoon Syn. ii. 114.

ORDO NAT. VERTICILLATÆ. Linn.

CLASSIS VIII. ORDO VI. LABIATÆ. III. Stamina quatuor fertilia. Corolla bilabiata. Calyx quinquefidus. Juss.

CHAR. GEN. Corollæ labium inferius tripartitum, laciniâ intermediâ subrenatâ. Stamina recta, distantia.

127. HYSSOPUS OFFICINALIS.

CHAR. SPEC. Floribus verticillatis racemosis secundis, laciniâ corollæ intermediâ bilobâ integerrima, foliis lineari-lanceolatis.

ANGLICE, Hyssop.

LOCUS. In Siberiæ, et Pannoniæ ruderatis, et montosis. Perennis.

VIS. Amara, tonica, stimulâs.

USUS. In catarrho, hysteriâ.

MENTHA. Willd. Spec. Plant. iii. 74. Perseon Syn. ii. 118.

ORDO NAT. VERTICILLATÆ. Linn.

CLASSIS VIII. ORDO VI. LABIATÆ. Juss.
Sectio III.

CHAR. GEN. Corolla subæqualis, quadrifida: laciniâ latiore emarginatâ. Stamina erecta distantia.

* Spicatæ.

128. MENTHA VIRIDIS.

CHAR. SPEC. Spicis oblongis interruptis, foliis lanceolatis nudis serratis sessilibus, staminibus corollâ longioribus. Tota glabra. Bracteæ setaceæ, flores longiores.

ANGLICE, Spear Mint.

LOCUS. In Germaniâ, Angliâ, Galliâ, Helvetiâ. Perennis.

VIS. Aromatica, calida. Stomachica, carminativa.

COMP. Aqua Menthæ viridis. *Lond. Eblan.* Inf. Menthæ Comp. *Eblan.* Oleum Menthæ viridis. *Lond.* et *Eblan.* Spiritus Menthæ viridis. *Lond.*

129. MENTHA PIPERITA.

CHAR. SPEC. Spicis obtusis inferne interruptis, foliis subovatis glabriusculis petiolatis, calycè basi glaberrimo.

ANGLICE, Peppermint.

LOCUS. Angliæ indigena. Perennis.

VIS. Eadem que præcedentis, sed acrior, cum frigoris sensû. Carminativa, stomachica.

USUS. Ad nauseam, flatum, et tormina amovenda.

COMP. Aqua Menthæ Pip. *Lond. Edin. et Eblan.*
Oleum Menthæ Pip. *Lond. Edin. et Eblan.* Spiritus
Menth. Pip. *Lond. et Edin.*

*** Verticillatæ.

130. MENTHA PULEGIUM.

CHAR. SPEC. Floribus verticillatis, foliis ovatis obtusis subcrenatis, caulibus subteretibus repentibus, staminibus corollâ longioribus.

ANGLICE, Pennyroyal.

LOCUS. In Germaniæ, Angliæ, Galliæ subhumidis. Perennis.

VIS. Fragrans, pungens, olim emmenagoga, expectorans, diaphoretica, existimata.

COMP. Aqua Pulegii. *Lond. Edin. et Eblan.* Oleum Pulegii. *Lond. et Eblan.* Spiritus Pulegii. *Lond.*

LAVANDULA. Willd. Spec. Plant. iii. 60. Persoon Syn. ii. 116.

ORDO NAT. VERTICILLATÆ. Linn.

CLASSIS VIII. ORDO VI. LABIATÆ. Sectio III. Juss.

CHAR. GEN. Calyx ovatus, subdentatus, bractea suffultus. Corolla resupinata. Stamina intra tubum.

131. LAVANDULA SPICA.

CHAR. SPEC. Foliis sessilibus lanceolato-linearibus, margine revolutis, spicâ interruptâ, nudâ.

ANGLICE, Lavender.

LOCUS. In Europâ Australiori. Perennis.

VIS. Fragrans, amaricans, acris.

COMP. Oleum Lavandulæ. *Lond. Edin. et Eblan.*
Spiritus Lavandulæ. *Lond. Edin. et Eblan.* Spir. Lavand. Comp. *Lond. Edin. et Eblan.* Pulv. Asari Comp. *Edin. et Eblan.*

TEUCRIUM. Willd. Spec. Plant. iii. 13. Persoon Syn. ii. 110.

ORDO NAT. VERTICILLATÆ. Linn.

CLASSIS VIII. ORDO VI. LABIATÆ. II. Stamina quatuor fertilia. Corolla unilabiata, labio superiore subnullo. Juss.

CHAR. GEN. Corollæ labium superius nullum, ultra basin bipartitum. Stamina exserta.

132. TEUCRIUM MARUM.

CHAR. SPEC. Foliis integerrimis ovatis acutis petiolatis, subtus tomentosis, floribus racemosis secundis.

ANGLICE, Common Marum.

LOCUS. Valentia. Perennis.

VIS. Aromatica, amaricans, errhina.

COMP. Pulv. Asari Comp. *Edin. et Eblan.*

133. TEUCRIUM CHAMÆDRYS.

CHAR. SPEC. Foliis cuneiformi-ovatis incisis crenatis petiolatis, floribus ternis, caulibus procumbentibus subpilosus. Bractea serrata.

ANGLICE, Wall Germander.

LOCUS. Italia, Gallia, Anglia, Germania, Helvetia. Perennis.

VIS. Amara, tonica, stomachica, diuretica, emmenagoga, existimata.

MARRUBIUM. Willd. Spec. Plant. iii. 109. Persoon Syn. ii. 125.

ORDO NAT. VERTICILLATÆ. Linn.

CLASSIS VIII. ORDO VI. LABIATÆ. Sectio III. Juss.

CHAR. GEN. Calyx hypocrateriformis, rigidus, decemstriatus. Corollæ labium superius bifidum, lineare, rectum.

** Calycibus decem-dentatis.

134. MARRUBIUM VULGARE.

CHAR. SPEC. Foliis subrotundo-ovatis dentatis rugoso-venosis, calycinis dentibus setaceis uncinatis.

ANGLICE, Horehound.

LOCUS. Europæ ruderata, argillosa. Perennis.

VIS. Graveolens, amara, tonica, diuretica.

USUS. Olim in hysteriâ, asthmâ, catarrho.

**** Calyces bilabiati.**

ORIGANUM. Willd. Spec. Plant. iii. 132. Persoon Syn. ii. 128.

ORDO NAT. VERTICILLATÆ. Linn.

CLASSIS VIII. ORDO VI. LABIATÆ. IV. Stamina quatuor fertilia. Corolla bilabiata. Calyx bilabiatus. Juss.

CHAR. GEN. Strobilus tetragonus, spicatus calyces colligens. Corollæ labium superius erectum, planum; inferius tripartitum; laciniis æqualibus.

135. ORIGANUM VULGARE.

CHAR. SPEC. Spicis subrotundis paniculatis conglomeratis, bracteis calyce longioribus ovatis coloratis.

ANGLICE, Wild Marjoram.

LOCUS. Europæ Montes. Perennis.

VIS. Amara, subacris.

COMP. Oleum origani. *Lond. et Eblan.*

136. ORIGANUM MAJORANA.

CHAR. SPEC. Spicis subrotundis ternis compactis, pedunculatis, foliis petiolatis ellipticis obtusis glabriusculis, radice annuâ.

ANGLICE, Sweet Marjoram.

LOCUS. Lusitania, Palæstina. Annuâ.

VIS. Fragrans, aromatica.

COMP. Pulv. Asari. Comp. *Edin. et Eblan.*

MELISSA. Willd. Spec. Plant. iii. 146. Persoon Syn. ii. 132.

ORDO NAT. VERTICILLATÆ. Linn.

CLASSIS VIII. ORDO VI. LABIATÆ. Sectio IV.

Juss.

CHAR. GEN. Calyx aridus, supra planiusculus; labio superiore subfastigiato. Corollæ labium superius subfor-
nicatum, bifidum; labium inferius lobo medio cordato.

137. MELISSA OFFICINALIS.

CHAR. SPEC. Verticillis dimidiatis subsessilibus,
bracteis oblongis pedicellatis, foliis ovatis acutis serratis.

ANGLICE, Balm.

LOCUS. In montibus, et ad Europæ sepes australioris.
Perennis.

VIS. Suaveolens, aromatica, stomachica, diuretica.

CLASSIS XIV.

ORDO II.

DIDYNAMIA. ANGIOSPERMIA.

***** *Calyces quinquefidi.*

SCROPHULARIA. Willd. Spec. Plant. iii. 269. Persoon
Syn. ii. 160.

ORDO NAT. PERSONATÆ. Linn.

CLASSIS VIII. ORDO VII. SCROPHULARIÆ. I.

Stamina quatuor didynama. Juss.

CHAR. GEN. Calyx quinquefidus. Corolla subglo-
bosa, resupinata. Capsula bilocularis.

138. SCROPHULARIA NODOSA.

CHAR. SPEC. Foliis cordatis trinervatis, caule ob-
tusangulo.

ANGLICE, Figwort.

LOCUS. Succulentis, humidis. Perennis.

VIS. Fœtida, amara.

USUS. Olim contra Scrophulam unde nomen.

DIGITALIS. Willd. Spec. Plant. iii. 283. Persoon Syn. ii. 162.

ORDO NAT. LURIDÆ. Linn.

CLASSIS VIII. ORDO VII. SCROPHULARIÆ.

Sectio I. Juss.

CHAR. GEN. Calyx quinquepartitus. Corolla campanulata, ventricosa quinquefida. Capsula ovata, bilocularis. (Flores racemo unico.)

139. DIGITALIS PURPUREA.

CHAR. SPEC. Calycinis foliolis ovatis acutis, corollis obtusis: labio superiore integro, foliis ovato-lanceolatis rugosis.

ANGLICE, Foxglove.

LOCUS. In sylvaticis præsertim arenosis. Biennis.

VIS. Amara, nauseosa, subacris, maximè sedativa.

Intra sex horas pulsus primò duritie, deinde frequentia (scilicet a 130 ad 40) in sexagessimâ horæ parte imminutus est: sed exercitatione subitâ, vel etiam modicâ, iterum acceleratus; cor fortitèr palpitât, nausea, syncope, convulsiones, et mors interdum superveniunt. Ita ut benè cavendum, ne ultrâ modum adhibita sit, nam etsi vis systematis arterialis in morbis phlogisticis reprimatur, hoc medicamen venæsectionis officio nunquam fungi potest. Diuretica habebatur, ideoque largè et incautè antehac usurpata fuit; nonnunquam dubio, interdum periculoso, sæpè lethali eventu. Ammonia et Spiritus vini Gallici, et corporis positio recumbens, his molestiis, Digitalis usum sequentibus, optimè obviam ibunt.

Præscripta est in Morbis Phlogisticis, Phthisi, Hæmorrhagiis activis, Hydrope, et Hydrothoraci.

DOSIS. Pulv. gr. ss. ad gr. ij. Tinct. m. x. ad m. xx. Inf. ʒij. ad ʒss.

COMP. Decoct. Digitalis. *Eblan.* Inf. Digitalis. *Lond. et Edin.* Tinct. Digitalis. *Lond. Edin. et Eblan.*

CLASSIS XV.

ORDO I.

TETRADYNAMIA. SILICULOSA.

** *Silicula emarginata apice.*

COCHLEAREA. Willd. Spec. Plant. iii. 448. Persoon Syn. ii. 185.

ORDO NAT. SILIQUOSÆ. Linn.

CLASSIS XIII. ORDO III. CRUCIFERÆ. II.

Fructus siliculosus. Stylus unicus. Juss.

CHAR. GEN. Silicula turgida, rugosa, polysperma, bivalvis: valvulis gibbis obtusis. Dissepimentum subparallelum.

140. COCHLEAREA ARMORACIA.

CHAR. SPEC. Foliis radicalibus subrotundis, caulinis subsinuatis, siliculis globosis.

ANGLICE, Horse-radish.

LOCUS. Europæ paludes australioris. Perennis.

VIS. Acris, pungens, subdulcis. Stimulans, diuretica.

USUS. Rheumatismo, Hydrope, Scorbuto.

COMP. Infusum Armoraciæ Comp. Lond. Spiritus Armoraciæ Comp. Lond. Eblan.

CLASSIS XV.

ORDO II.

TETRADYNAMIA. SILIQUOSA.

** *Calyx hians foliolis superne distantibus.*

CARDAMINE. Willd. Spec. Plant. iii. 481. Persoon Syn. ii. 195.

ORDO NAT. SILIQUOSÆ. Linn.

CLASSIS XIII. ORDO III. CRUCIFERÆ. I.

Fructus siliquosus. Stylus nullus. Juss.

CHAR. GEN. Siliqua longa elastice dissiliens, valvulis revolutis, (in plurimis.) Stigma integrum. Calyx subhians. Glandula utriusque solitaria inter stamina breviora et calycem.

141. CARDAMINE PRATENSIS.

*** Foliis pinnatifidis, aut pinnatis.

CHAR. SPEC. Foliolis radicalibus subrotundis, caulinis lineari-lanceolatis.

ANGLICE, Lady's Smock.

LOCUS. In pascuis aquosis. Perennis.

VIS. Subacris, amaricans. Stimulans, diaphoretica?

USUS. Olim in spasmis.

SINAPIS. Willd. Spec. Plant. iii. 554. Persoon Syn. ii. 207.

ORDO NAT. SILIQUOSÆ. Linn.

CLASSIS XIII. ORDO III. CRUCIFERÆ. Sectio I. Juss.

CHAR. GEN. Calyx patens. Corollæ ungues recti. Glandula inter stamina breviora et pistillam, interque longiora et calycem. Dissepimentum prominens, ensiforme.

142. SINAPIS ALBA.

CHAR. SPEC. Siliquis hispidis subtetraspermis, rostro obliquo longissimo ensiformi. Pili in siliquis albid.

ANGLICE, White Mustard.

LOCUS. In agris indigena. Annua.

VIS. Pungens, amaricans. Semina oleum expressione effundunt. Stimulans, diuretica, emetica, rubefaciens.

USUS. In Dyspepsiâ, Chlorosi, et alvi obstipatione seminum integrorum cochleare medium, bis terve die ingurgitatur.

Periculosa tamen eorum usum subindè sequuta sunt, nempe Stomachi inflammatio etiam lethalis, sive Enteritis, seminibus intra Cæci appendicem vermiformem transit. Per

bettoniadas etiam septem intra primas vias semina retineri cognovimus.

Morbis caput afficientibus, cataplasma e seminum pulvere confectum, cuti quam citissime rubefaciendas subservit.

COMP. Cataplasma Sinapis. *Lond. et Eblan.*

143. SINAPIS NIGRA.

CHAR. SPEC. Siliquis glabris tetragonis sublævibus, racemo adpressis, foliis summis lineari-lanceolatis. Rostrum quadrangulum.

ANGLICE, Black Mustard.

LOCUS. Indigena, aggeribus ruderatis. Annuæ.

CLASSIS XVI.

ORDO I.

MONADELPHIA. TRIANDRIA.

TAMARINDUS. Willd. Spec. Plant. iii. 577. Persoon Syn. ii. 211.

ORDO NAT. LOMENTACEÆ. Linn.

CLASSIS XIV. ORDO XI. LEGUMINOSÆ. I.

Corolla regularis. Legumen multiloculare, sæpius bivalve, dissepimentis transversis, loculis monospermis. Stamina distincta. Arbores aut frutices, folia abruptè pinnata. Juss.

CHAR. GEN. Calyx quadripartitus, basi turbinatus. Petala tria, ascendentia. Filamenta basi tantùm coalita; tria longiora fertilia, quatuor minima sterilia fertilibus interposita, duo setiformia sterilia priorum vaginæ extus opposita. Legumen duplici cortice vestitum: exteriore sicco fragili, interiore membranaceo, in utriusque interstitio pulposum, uni-vel triloculare.

144. TAMARINDUS INDICA.

ANGLICE, Tamarind.

LOCUS. In Indiâ, Americâ, Ægypto, Arabiâ. Perennis.

VIS. Acido-dulcis, laxativa, refrigerans in febris.

COMP. Inf. Tamarindi cum Sennâ. *Edin. et Eblan.*
Conf. Sennæ.

CLASSIS XVI.

ORDO VIII.

MONADELPHIA. POLYANDRIA.

MALVA. Willd. Spec. Plant. iii. 774. Persoon Syn. ii. 249.

ORDO NAT. COLUMNIFERÆ. Linn.

CLASSIS XIII. ORDO XIV. MALVACEÆ. II. Stamina in tubum corolliferum connata, indefinita. Fructus multicapsularis, capsulæ verticillatæ, in orbem dispositæ, aut in unam compactæ. Juss.

CHAR. GEN. Calyx duplex; exterior triphyllus. Capsulæ plurimæ monospermæ.

** Foliis angulatis.

145. **MALVA SYLVESTRIS.**

CHAR. SPEC. Caule erecto herbaceo, foliis septemlobatis acutis, pedunculis petiolisque pilosis.

ANGLICE, Common Mallow.

LOCUS. Europæ campestribus. Biennis.

VIS. Fatua, mucilaginosa.

USUS. Olim Dysenteriâ, Ischuriâ.

ALTHÆA. Willd. Spec. Plant. iii. 770. Persoon Syn. ii. 248

ORDO NAT. COLUMNIFERÆ. Linn.

CLASSIS XIII. ORDO XIV. MALVACEÆ. Sectio II. Juss.

CHAR. GEN. Calyx duplex: exterior sex-sive novemfidus. Capsulæ plurimæ monospermæ.

* Calyce exteriore octofido, sive novemfido. Capsulis emarginatis.

146. ALTHÆA OFFICINALIS.

CHAR. SPEC. Foliis tomentosis oblongo-ovatis obsoletè trilobis dentatis.

ANGLICE, Marsh Mallow.

LOCUS. Sponte in Belgiæ, Angliæ, Galliæ, Siberiæ humidis. Perennis.

VIS. Mucilaginosa, subdulcis.

USUS. In Pulmonum, et intestinorum morbis olim usurpata.

COMP. Decoctum Althææ Officinalis. *Edin.* *Syr.* Althææ. *Lond.* et *Edin.*

CLASSIS XVII.

ORDO III.

DIADELPHIA. OCTANDRIA.

POLYGALA. Willd. Spec. Plant. iii. 871. Persoon Syn. ii. 271.

ORDO NAT. VAGÆ. Linn.

CLASSIS VIII. ORDO II. PEDICULARES. Sectio I. Juss.

CHAR. GEN. Calyx pentaphyllus: foliolis duobus alæformibus, coloratis. Capsula obcordata, bilocularis, bivalvis.

*** Imberbes, herbacææ.

147. POLYGALA SENEGA.

CHAR. SPEC. Floribus imberbibus, spica terminali filiformi, caule erecto herbaceo simplicissimo, foliis oblongo-lanceolatis.

ANGLICE, Senega.

LOCUS. Americâ Boreali. Perennis.

VIS. Acri-acida, subaromatica, stimulans, expectorans?

USUS. In Peripneumoniâ, Rheumatismo.

DOSIS. ʒss. ad ʒii,

COMP. Dec. Senegæ. Lond. et Eblan.

CLASSIS XVII.

ORDO IV.

DIADELPHIA. DECANDRIA.

* *Stamina omnia connexa.*

PTEROCARPUS. Willd, Spec. Plant. iii. 904. Persoon Syn. ii. 277.

CLASSIS XIV. ORDO XI. LEGUMINOSÆ. IX.
Corolla irregularis papilionacea. Stamina plerumque decem diadelpha. Legumen capsulare uniloculare submonospermum sæpè non dehiscens. Arbores aut frutices, folia sæpius impari-pinnata; stipulæ a petiolo distinctæ mox deciduæ. Juss.

CHAR. GEN. Calyx quinquedentatus. Legumen orbiculari-falcatum, foliaceum inæqui-alatum non dehiscens. Semina aliquot solitaria.

148. PTEROCARPUS SANTALINUS.

CHAR. SPEC. Foliis ternatis subrotundis retusis glaberrimis, petalis crenatis, undulatis.

ANGLICE, Red Saunders.

LOCUS. Indiæ Orientalis montibus, Zeylonâ. Perennis.

VIS. Suaveolens, rubeo-tinctoria.

149. PTEROCARPUS ERINACEA.

CHAR. SPEC. Foliis pinnatis, foliolis oblongis tomentoso-canescens, fructibus medio piloso-hispidis.

ANGLICE, Kino.

LOCUS. Senegallia.

VIS. Dalcamara, rubicundo-fusca, styptica: continet Tannin, et acidum Gallicum.

USUS. In Diarrhœa, Hæmorrhagiis passivis, et Leucorrhœa.

DOSIS. Gr. x. ad ʒi. Pulv.

COMP. Tinct. Kino. *Lond. Edin. et Eblan.* Elect. Catechu. *Edin. et Eblan.*

Eucalyptus Resinifera. Edin. Butea Frondosa. Eblan.

SPARTIUM. Willd. Spec. Plant. iii. 926. Persoon Syn. ii, 286.

ORDO NAT. PAPILIONACEÆ. Linn.

CLASSIS XIV. ORDO XI. LEGUMINOSÆ. V.

Corolla irregularis papilionacea. Stamina decem diadelphea. Legumen uniloculare bivalve. Frutices aut herbæ; folia simplicia, aut ternata aut rarius digitata; stipulæ nunc subnullæ, nunc conspicuæ imo petiolo adnatæ, aut ab eodem distinctæ. Juss.

CHAR. GEN. Calyx deorsum productus. Carina quibusdam pendula. Filamenta germini adhærentia. Stigma longitudinale suprâ villosum. Legumen oblongum, mono- aut poly-spermum.

**** Foliis ternatis.**

150. SPARTIUM SCOPARIUM.

CHAR. SPEC. Foliis ternatis solitariisque oblongis, floribus axillaribus, leguminibus margine pilosis, ramis angulatis.

ANGLICE, Broom.

LOCUS. Indigena, in arenosis. Perennis.

VIS. Amara, diuretica præstans, laxativa.

USUS. Hydropes.

DOSIS. ʒi ad ʒi.

COMP. Ext. Cacuminum Genistæ. *Eblan.*

**** Stigma pubescens.** (Nec priorum notæ.)

DOLICHOS. Willd. Spec. Plant. iii. 1037.

STIZOLOBIUM. Persoon Syn. ii. 298.

ORDO NAT. PAPILIONACEÆ. Linn.

CLASSIS XIV. ORDO XI. LEGUMINOSÆ. Sectio V. Juss.

CHAR GEN. Calyx campanulatus, bilabiatus; labio superiore integro-erecto; inferiore trifido: lacinia media productiore vexillum assurgens. Alæ dolabræformes basi lunatæ, longitudine carinæ. Antheræ biformes, hirsutæ, legumen torosum, uniloculare: dissepimentis interceptum. Semina orbicularia, hilo cristata ultra medium cincta.

151. STIZOLOBIUM PRURIENS.

CHAR. SPEC. Volubilis leguminibus racemosis: valvulis subcarinatis hirtis, pedunculis-ternis.

STIZOLOBIUM PRURIENS. Persoon Syn. ii. 299.

CHAR. SPEC. Leguminibus racemosis: valvulis subcarinatis hirtis, pedunculis ternis.

ANGLICE, Cowhage.

LOCUS. America, India Orientalis, et Occidentalis. Perennis.

VIS. Leguminum Setæ pruritum intolerandum excitant. Anthelmintica mechanica, contra lumbricos teretes, et ascariides: sed haud incautè dari oportet, quia interdum Enteritin, aliàs pruritum molestissimum circa anum, inducunt.

DOSIS. Setarum. Gr. v. ad gr. x.

COMP. Confectio Dolichos Prurientis.

***** Legumen subbiloculare.** (Nec Priorum.)

ASTRAGALUS. Willd. Spec. Plant. iii. 1256. Persoon Syn. ii. 334.

ORDO NAT. PAPILIONACEÆ. Linn.

CLASSIS XIV. ORDO XI. LEGUMINOSÆ.

VI. Corolla irregularis papilionacea. Stamina decem diadelpa. Legumen uniloculare bivalve. Herbæ, aut frutices, aut arbusculæ; folia impari pinnata. Juss.

CHAR. GEN. Carina obtusa, legumen biloculare, aut subbiloculare. Sutura inferiore introflexa.

Floribus pedunculatis.

152. ASTRAGALUS TRAGACANTHA.

CHAR. SPEC. Frutescens, petiolis spinescentibus, foliolis ellipticis incanis, pedunculis subquadrifloris, folia æquantibus calycinis dentibus ovatis brevibus.

Nullum gummi fluit ex hac planta. Persoon Syn. ii. 341.

• ASTRAGALUS CRETICUS.

CHAR. SPEC. Frutescens, petiolis spinescentibus, foliis tri-quadri-jugisve, foliolis lanceolatis incanis, floribus axillaribus sessilibus solitariis calyce lanato, caulis nigricans, corolla lineis purpureis striata.

Ex hoc frutice legitur gummi Tragacanthæ album, ut ex Itinerario Tournefortii patet. Reliquæ species, excepto Astragalo gummifero, qui gummi flavescens minoris pretii fundit, nullo scatent gummoso. Persoon Syn. ibid.

ANGLICE, Tragacanth.

LOCUS. Creta Insula; ad montem Ida. Perennis.

VIS. Mucilaginis.

COMP. Mucilago Astragalæ Tragacanthæ. *Edin.* et *Eblan.* Pulv. Tragacanthæ Comp. *Lond.*

**** Legumina Submonosperma. (Nec Priorum.)

GLYCYRRHIZA. Willd. Spec. Plant. iii. 1143.

ORDO NAT. PAPILIONACEÆ. Linn.

CLASSIS XIV. ORDO XI. LEGUMINOSÆ.

Sectio VI. Juss.

LIQUIRITIA. Persoon Syn. ii. 313.

CHAR. GEN. Calyx tubulosus æqualis, quinquepartitus, corollæ vexillo erecto, lateribus reflexo, alis patentibus, carina bifida. Legumen oblongum glabrum, tri-vel tetraspermum, (folia exstipulata flores racemosi.)

153. LIQUIRITIA OFFICINALIS.

ANGLICE, Liqueurice.

LOCUS. In Europa Australiore. Perennis.

VIS. Dulcis mucilaginosa, aloen, et alia corrigens.

COMP. Dec. Sarsaparillæ Comp. *Lond. Eblan.* Infusum Lini. Comp. *Lond.* Ext. Glycyrrhizæ. *Lond. Edin.* et *Eblan.* Confectio Sennæ. *Lond. Edin.*

*****Legumen uniloculare, polyspermum. (Nec priorum.)

GEOFFRÆA. Willd. Spec. Plant. iii. 1129. Persoon Syn. ii. 278.

CLASSIS XIV. ORDO XI. LEGUMINOSÆ.

Sectio IX. Juss.

CHAR. GEN. Calyx campanulatus, semiquinquefidus subulabatus. Alæ et carina subæquales; vexillum majus. Drupa ovoidea, utrinque sulco notata. Nux bivalvis, monosperma.

154. GEOFFRÆA INERMIS.

CHAR. GEN. Inermis, foliolis ovato-lanceolatis. Folia subpedalia.

ANGLICE, Cabbage-tree.

LOCUS. In Jamaicae sylvis. Perennis.

VIS. Subdulcis, mucilaginosa, anthelmintica, cathartica.

DOSIS. Pulv. ℞. ad iii.

COMP. Decoctum Geoffrææ Inermis. *Edin.*

CLASSIS XVIII.

ORDO III.

POLYADELPHIA. ICOSANDRIA.

CITRUS. Willd. Spec. Plant. iii. 1426. Persoon Syn. ii. 73.

ORDO NAT. HESPERIDEÆ. Linn.

CLASSIS XHI. ORDO X. AURANTIA. II. Fructus polyspermus baccatus. Folia pellucido-punctata. Aurantia vera. Juss.

CHAR. GEN. Calyx quinquefidus. Petala quinque, oblonga. Filamenta dilatata, basi subconnata in fasciculos, bacca corticosa, 9-18-ocularis. Pulpa molli.

155. CITRUS MEDICA.

CHAR. SPEC. Petiolis linearibus, foliis ovatis acuminatis.

ANGLICE, Lemon.

LOCUS. Asiâ, et Europâ Meridionali. Perennis.

VIS. Cortex fragrans, odoratus, vesiculis oleosis, aromatica, amara. Pulpa acida. Succus refrigerans, antisepticus. Acidum citricum, mucilaginem, saccharum et aquam continet.

USUS. In Febribus, et Scorbuto.

COMP. Acidum Citricum. *Lond.* et *Eblan.* *Syr.* Limonis. *Lond.* et *Edin.*

156. CITRUS AURANTIUM.

CHAR. SPEC. Petiolis alatis, foliis ovato-lanceolatis, acuminatis. Fructus subglobosus.

ANGLICE, Orange.

LOCUS. In Indiâ, et ubique nunc colitur. Perennis.

MELALEUCA Willd. Spec. Plant. iii. 1428. Persoon Syn. ii. 26.

CLASSIS XIV. ORDO VII. MYRTL. I. Flores in foliorum axillis, aut in pedunculis multifloris oppositi. Folia plerumque opposita et punctata. Juss.

CHAR. GEN. Calyx quinquepartitus, semisuperus. Petala quinque. Stamina in quinque fasciculos, connata sive ramosa. Capsula infera semivestita, trilocularis. Semina immatura lineari-paleacea, matura sæpius alata.

* Foliis alternis.

157. MELALEUCA LEUCODENDRON.

MELALEUCA CAJUPUTI. Phar. Lond.

CHAR. SPEC. Foliis alternis lanceolatis, acuminatis falcato-obliquis quinque-nerviis, ramulis petiolisque glabris. Arbor trunco nigro ramis albis, ex quâ verum Oleum Cajeputi obtinetur.

LOCUS. India Orientalis. Perennis.

VIS. Stimulans, antispasmodica, diaphoretica.

CLASSIS XIX.

ORDO I.

SYNGENESIA. ÆQUALIS.

* *Semiflosculosi, corollis ligulatis omnibus.*

LEONTODON. Willd. Spec. Plant. iii. 1544. Persoon Syn. ii. 367.

ORDO NAT. COMPOSITI a. SEMIFLOSCULOSI.
Linn.

CLASSIS X. PLANTÆ DICOTYLEDONES MONOPETALÆ. COROLLA EPIGYNA. ANTHERÆ CONNATÆ. ORDO I. CICHORACEÆ. III. Receptaculum nudum. Semen papposum pappo plumoso. Juss.

CHAR. GEN. Calyx duplex. Receptaculum nudum. Pappus stipitatus, plumosus.

158. LEONTODON TARAXACUM.

CHAR. SPEC. Calyce exteriori reflexo, scapo unifloro, foliis runcinatis glabris, laciniis lanceolatis dentatis.

ANGLICE, Dandelion,

LOCUS. Europæ pascua aprica. Perennis.

VIS. Amariçans, tonica in Dyspepsiâ.

LACTUCA. Willd. Spec. Plant. iii. 1523. Persoon Syn. ii. 364.

ORDO NAT. COMPOSITI a. SEMIFLOSCULOSI.
Linn.

CLASSIS X. ORDO I. CICHORACEÆ. II. Re-
ceptaculum nudum. Semen papposum pappo piloso. Juss.

CHAR. GEN. Calyx imbricatus, cylindricus, margine
membranaceo. Receptaculum nudum. Pappus simplex,
stipitatus. Semina lævia.

159. LACTUCA SATIVA.

CHAR. SPEC. Foliis rotundatis, caulinis, cordatis,
caule corymboso.

ANGLICE, Lettuce.

LOCUS. In Europâ. Annua.

VIS. Lactescens, narcotica.

Folia, et caulis succo proprio gaudent; qui florescente
plantâ, ex caule inciso defluit, aëre spissatus, Lactucarium
appellatur; et, Morphiam alkali continendo, ad opii simili-
tudinem accedit. Ad Phthiseos tussim sublevandam nuper
datum est.

DOSIS. Lactucarii gr. i. ad vi.

COMP. Succus spissatus Lactusæ Sativæ. *Edin.*

160. LACTUCA VIROSA.

CHAR. SPEC. Foliis horizontalibus, carina aculeatis
denticulatis, apice obtusis basi sagittatis, oblongo-lanceolatis,
infimis sinuatis.

ANGLICE, Opium-scented Lettuce.

LOCUS. In Europæ aggeribus Australioris. Annua.

VIS. Graveolens, amara, narcotica, venenata.

COMP. Succus spissatus Lactusæ virosæ. *Edin.*

**** Capitati.**

ARCTIUM. Willd. Spec. Plant. iii. 1631. Persoon
Syn. ii. 383.

ORDO NAT. COMPOSITI. b. CAPITATI. Linn.

CLASSIS X. ORDO II. CINAROCEPHALÆ. I.
Cinarocephalæ veræ. Squamæ calycis spinosæ. Juss.

CHAR. GEN. Calyx globosus, squamis apice hamis
inflexis. Receptaculum paleaceum. Pappus setoso-pa-
leaceus.

161. ARCTIUM LAPPA.

CHAR. SPEC. Foliis caulinis cordatis, petiolatis, denticulatis; calycibus glabris.

ANGLICE, Burdock.

LOCUS. Europæ indigena. Biennis.

VIS. Amara, diuretica.

USUS. Olim Scorbuto, Arthritide.

CLASSIS XIX.**ORDO II.****SYNGENESIA. SUPERFLUA.******* Discoidei.**

ARTEMISIA. Willd. Spec. Plant. iii. 1815. Persoon Syn. ii. 409.

ORDO NAT. NUCAMENTACEÆ. Linn.

CLASSIS X. ORDO III. CORYMBIFERÆ. IV.
Receptaculum nudum. Semen nudum seu non papposum.
Flores flosculi. Juss.

CHAR. GEN. Calyx imbricatus, squamis retundatis, conniventibus. Corollæ radii nulli. Receptaculum subvillosum, vel nudiusculum. Pappus nullus. (Flores ut plurimum racemosi.)

* Fruticosæ.

162 ARTEMISIA ABROTANUM.

CHAR. SPEC. Frutescens, caule stricto, foliis inferioribus bipinnatis, superioribus pinnatis capillaceis, calycibus pubescentibus hemisphæricis.

ANGLICE, Southernwood.

LOCUS. Siberiâ, Cappadaciâ, Syriâ, Galatiâ. Perennis.

VIS. Amara, tonica.

USUS. Olim pro fomentis, Gangrænâ.

*** Herbaceæ, caule ramoso, paniculato, foliis compositis.

163. ARTEMISIA SANTONICA.

CHAR. SPEC. Foliis caulinis pinnatis linearibus glabris, ramis indivisis, spicis secundis reflexis, floribus subsessilibus quinquefloris.

ANGLICE, Tartarian Southernwood.

LOCUS. In Tartariâ, Persiâ. Perennis.

VIS et USUS. Præcedentis, A. Abrotani.

164. ARTEMISIA MARITIMA.

CHAR. SPEC. Foliis niveo-tomentosis, caulinis, pinnatis, linearibus, obtusis, rameis, linearibus simplicibus obtusis, caule ascendente ramoso, ramulis cernuis, floribus oblongis tomentosis sessilibus.

ANGLICE, Sea Wormwood.

LOCUS. In Norvegiâ, Pomeraniæ, Belgii, Angliæ maritimæ. Perennis.

VIS. Sequentis, A. Absinthii.

165. ARTEMISIA ABSINTHIUM.

CHAR. SPEC. Foliis incanis: radicalibus triplicatopinnatifidis laciniis lanceolatis dentatis obtusis, caulinis bipinnatifidis, pinnatifidisve, laciniis lanceolatis acutiusculis, floralibus indivisis lanceolatis, floribus globosis pedunculatis, nutantibus.

ANGLICE, Common Wormwood.

LOCUS. Europæ ruderata. Perennis.

VIS. Amarissima, tonica, anthelmintica.

COMP. Ext. Absinthii, *Phar. Eblan.*

TANACETUM. Willd. Spec. Plant. iii. 1809. Persoon Syn. ii. 498.

ORDO NAT. COMPOSITI. c. CORYMBIFERI.
Lin.

CLASSIS X. ORDO III. CORYMBIFERÆ. Sectio IV. Juss.

CHAR. GEN. Calyx imbricatus, hemisphæricus, squamis acuminatis. Corollæ radii obsoletè trifidi. Receptaculum nudum. Pappus submarginatus.

166. TANACETUM VULGARE.

CHAR. SPEC. Foliis bipinnatis incisiss serratis.

ANGLICE, Tansy.

LOCUS. Europæ margines pratorum. Perennis.

VIS. Amara, aromatica.

USUS. Olim Hydrope, Hysteria, &c.

*** Radiati.

ARNICA. Willd. Spec. Plant. iii. 2106. Persoon Syn. ii. 453.

CLASSIS X. ORDO III. CORYMBIFERÆ. II. Receptaculum nudum. Semen papposum. Flores radiati. Juss.

CHAR. GEN. Calyx foliolis æqualibus. Corollæ radii sæpiùs filamentis quinque absque antheris. Receptaculum nudum. Pappus simplex.

167. ARNICA MONTANA.

CHAR. SPEC. Foliis ovatis integris: caulinis geminis oppositis.

ANGLICE, Mountain Arnica.

LOCUS. In Europâ, et Americâ Septentrionalibus. Perennis.

VIS. Nauseosa, stimulans, emetica, cathartica.

USUS. Olim in Paralysi, Arthritide, &c.

INULA. Willd. Spec. Plant. iii. 2089. Persoon Syn. ii. 450.

ORDO NAT. COMPOSITI. c. CORYMBIFERI. Linn.

CLASSIS X. ORDO III. CORYMBIFERÆ. Sectio II. Juss.

CHAR. GEN. Calyx squarrosus, aut imbricatus, corollæ

radii-numerosi, lutei. Antheræ basi in setas duas desinentes. Receptaculum nudum. Pappus simplex.

168. INULA HELENIUM.

CHAR. SPEC. Foliis amplexicaulibus ovatis rugosis : subtus tomentosis. Calycum squamis ovatis.

ANGLICE, Elecampane.

LOCUS. In Europâ Australiori. Perennis.

VIS. Aromatica, mucilaginis.

USUS. Olim Dyspepsiâ, &c.

SOLIDAGO. Willd. Spec. Plant. iii. 2053. Persoon Syn. ii. 447.

ORDO NAT. COMPOSITI. c. CORYMBIFERI. Linn.

CLASSIS X. ORDO III. CORYMBIFERÆ. Sectio II. Juss.

CHAR. GEN. Calycis squamæ imbricatæ, clausæ. Corollæ radii circiter quinque (flavi.) Receptaculum nudum, punctis exaratum. Pappus simplex.

*** Racemis erectis.

169. SOLIDAGO VIRGAUREA.

CHAR. SPEC. Caule erecto tereti pubescente supernè ramoso, foliis inferioribus ellipticis pilosiusculis acutè serratis, racemis lateralibus et terminalibus erectis, ligulis elongatis. Caule apice subcompressus, et ibi magis pubescens. Radius octapetalus.

ANGLICE, Golden Rod.

LOCUS. In pascuis siccis, et sylvis. Perennis.

VIS. Astringens.

USUS. Lithontriptica olim existimata.

TUSSILAGO. Willd. Spec. Plant. iii. 1962. Persoon Syn. ii. 455.

ORDO NAT. COMPOSITI. c. CORYMBIFERI. Linn.

CLASSIS X. ORDO III. CORYMBIFERÆ. Sectio I. Juss.

CHAR. GEN. Calyx simplex, squamæ æquales, discum æquantēs, submembranaceæ. Corollæ fœminæ ligulatæ sive edentulæ. Receptaculum nudum. Pappus simplex, sessilis. Flosculi plerisque polygamo-dioici.

170. TUSSILAGO FARFARA.

CHAR. SPEC. Scapo unifloro subnudo bracteato, flore radiato, foliis cordatis angulatis dentatis subtus pubescentibus.

ANGLICE, Coltsfoot.

LOCUS. Europæ argillosa. Perennis.

VIS. Subacris, expectorans?

USUS. Olim in Catarrho, &c.

ANTHEMIS. Willd. Spec. Plant. iii. 2174. Persoon Syn. ii. 465.

ORDO NAT. COMPOSITI. c. CORYMBIFERI.
Linn.

CLASSIS X. ORDO III. CORYMBIFERÆ. V. Receptaculum paleaceum. Semen nudum seu non papposum. Flores plerumque radiati raro flosculosi. Juss.

CHAR. GEN. Calyx hemisphæricus, subæqualis. Flosculorum radii plures quam quinque. Receptaculum paleaceum: paleis planis apice acuminatis rigidis. Pappus nullus, sive margo membranaceus.

* Radio discolore sive albo.

171. ANTHEMIS NOBILIS.

CHAR. GEN. Foliis bipinnatis, foliolis tripartitis, lineari-subulatis subvillosis, caule basi ramoso.

ANGLICE, Chamomile.

LOCUS. Europæ pascua aprica. Annua.

VIS. Amara, suaveolens.

USUS. In Dyspepsiâ, &c. et pro fomentis.

DOSIS. Pulv. Florum ʒss. ad ʒi.

COMP. Dec. Anthemidis Nobilis. *Edin.* et *Hblan.*
Inf. Anthem. *Lond.* Ext. Anthem. *Lond.* et *Edin.* Oleum Anthemidis. *Lond.*

172. ANTHEMIS PYRETHRUM.

CHAR. SPEC. Foliis triplicato-pinnatis, foliolis linearibus, caule decumbente. Ramis axillaribus unifloris. Radius albus subtus purpureus. Radix crassa, sapore fervido, salivam excitat.

ANGLICE, Pellitory of Spain.

LOCUS. In Europâ Australiori. Perennis.

VIS. Acris, Sialogoga.

USUS. Odontalgia, &c.

CLASSIS XIX.**ORDO III.****SYNGENESIA. FRUSTRANEA.**

CENTAUREA. Willd. Spec. Plant. iii. 2277. Persoon Syn. ii. 481.

ORDO NAT. COMPOSITI. b. CAPITATI. Linn.

CLASSIS X. ORDO II. CINAROCEPHALÆ. II. Cinarocephalæ veræ. Squamæ calycis inermes. Plantæ plerumque inermes. Juss.

CHAR. GEN. Calyx varius. Corollæ radii infundibuliformes, longiores, irregulares. Receptaculum setosum. Pappus simplex, varius.

***** Calcitrapa. Calycis squamis spina solitaria, dentata, ramosa magna terminali.

173. CENTAUREA BENEDICTA.

CHAR. SPEC. Calycibus ramoso spinosis involucri lanatis, foliis semidecurrentibus, denticulato-spinosis.

ANGLICE, Blessed Thistle.

LOCUS. In Chio, Lemno, et Europâ Australiori. **Annua.**

VIS. Amarissima, emetica, diaphoretica.

USUS. In Dyspepsiâ, &c.

CLASSIS XX.

ORDO IV.

GYNANDRIA. HEXANDRIA.

ARISTOLOCHIA. Willd. Spec. Plant. iv. 151. Persoon Syn. ii. 526.

ORDO NAT. SARMENTACEÆ. Linn.

CLASSIS V. ORDO I. ARISTOLOCHIÆ. Juss.

CHAR. GEN. Calyx nullus. Corolla monopetala, ligulata, basi ventricosa. Capsula sexlocularis polysperma, infera.

** Caule debili, prostrato.

174. ARISTOLOCHIA SERPENTARIA.

CHAR. SPEC. Foliis cordatis, oblongis acuminatis, caule flexuoso ascendente, pedunculis radicalibus. Corollæ labio lanceolato.

ANGLICE, Virginian Snake Root, or Birthwort.

LOCUS. In Virginia. Perennis.

VIS. Amara, stimulans, diaphoretica.

DOSIS. Radicis pulveris gr. x. ad 3ss.

COMP. Tinct. Serpentariæ. Lond. Edin. et Eblan. Tinct. Cinchonæ Comp. Lond. et Eblan. Electuarium Opiatum. Edin.

CLASSIS XXI.

ORDO I.

MONGECIA. MONANDRIA.

EUPHORBIA. Willd. Spec. Plant. ii. 881. Persoon Syn. ii. 10.

ORDO NAT. TRICOCCEÆ. Linn.

CLASSIS XV. ORDO I. EUPHORBIÆ. I. Styli
plures definiti, sæpius tres. Juss.

CHAR. GEN. Calyx monophyllus, ventricostus. Corolla tetra-sive penta-petala calyci insidens. Capsula trilocca. (Plantæ lactescentes.) Stamina diverso tempore assurgentia.

* Aculeata.

175. EUPHORBIA OFFICINARUM.

CHAR. SPEC. Aculeata nuda multangularis: aculeis geminatis.

ANGLICE, Official Euphorbia.

LOCUS. In Æthiopiâ et Africâ calidlore. Perennis.

VIS. Potentissimè corrodens, rubefaciens, vesicans, ex-corians.

USUS. Olim in Hydrope, Paralyti, Amaurosi.

CLASSIS XXI.

ORDO IV.

MONECIA. TETRANDRIA.

MORUS. Willd. Spec. Plant. iv. 368. Persoon Syn. ii. 557.

ORDO NAT. SCABRIDÆ. Linn.

CLASSIS XV. ORDO III. URTICÆ. II. Flores receptaculo communi multifloro impositi, aut squamis involu-crantibus capitati aut distincti sparsi. Juss.

CHAR. GEN. Masculus. Calyx quadripartitus. Corolla nulla. Fœmineus. Calyx tetraphyllus. Corolla nulla. Styli duo. Calyx baccatus. Semen unicum.

176. MORUS NIGRA.

CHAR. SPEC. Foliis cordatis ovatis, aut subquinque-lobis inæqualiter dentatis scabris.

ANGLICE, Mulberry.

LOCUS. In Chinâ, et Italiâ. Perennis.

VIS. Acido-dulcis, rubro-tinctoria.

COMP. Syrupus Mori. *Phar. Lond.*

CLASSIS XXI.

ORDO VI.

MONŒCIA. HEXANDRIA.

COCOS. Willd. Spec. Plant. iv. 400. Persoon Syn. ii. 562.

CLASSIS III. PLANTÆ MONOCOTYLEDONES.
STAMINA PERIGYNA. ORDO I. PALMÆ. I.

Frondes pinnatæ foliola sæpius lanceolata angusta. Juss.

CHAR. GEN. Palma. Masculus. Calyx triphyllus. Corolla tripetala. Fœmineus. Calyx diphyllus. Corolla hexipetala. Stylus nullus. Stigma fovea. Drupa fibrosa.

177. COCOS BUTYRACEA.

CHAR. SPEC. Inermis, frondibus pinnatis, foliolis simplicibus.

ANGLICE, Maccaw tree.

LOCUS. Brasilia. Perennis.

Nuces contusæ, et in aquam calidam conjectæ, oleum flavum, suaveolens butyraceum suppeditant, quod vulgo dicitur *Palm Oil*.

CLASSIS XXI.

ORDO VII.

MONŒCIA. POLYANDRIA.

ARUM. Willd. Spec. Plant. iv. 477. Persoon Syn. ii. 574.

ORDO NAT. PIPERITÆ. Linn.

CLASSIS II. ORDO I. AROIDEÆ. I. Spadix spathâ involutus. Juss.

CHAR. GEN. Spatha monophylla, cucullata. Spadix suprâ nudus, infernè: fœmineus, medio stamineus. Infra glandulæ cirrhosæ. Bacca ut plurimum monosperma.

** Foliis simplicibus.

178. ARUM MACULATUM.

CHAR. SPEC. Foliis hastato-sagittatis, lobis deflexis, spadice clavato spathâ brevior. Flores fusco-maculati.

ANGLICE, Cuckoo-pint.

LOCUS. In Europæ sylvaticis. Perennis.

VIS. Acris, rubefaciens.

USUS. Olim in Cachexiâ, Chlorosi, Rheumatismo, &c.

QUERCUS. Willd. Spec. Plant. iv. 423: Persoon Syn. ii. 567.

ORDO NAT. AMENTACEÆ. Linn.

CLASSIS XV. ORDO IV. AMENTACEÆ. III. Flores monoici. Juss.

CHAR. GEN. Masculus. Calyx subquinquefidus. Stamina quinque ad decem. Fœmineus. Calyx monophyllus, integerrimus, scaber. Styli duo ad quinque. Nux coriacea, calyce persistente basi cincta.

** Foliis dentatis.

179. QUERCUS INFECTORIA.

CHAR. SPEC. Foliis oblongis mucronato-dentatis utrinque glabris.

ANGLICE, Dyers' Oak.

LOCUS. Asia Minor, a Bosphoro ad Syriam usque. Perennis.

Ex ramusculorum hujusce arboris surculis enascuntur tabera, Gallæ dicta, insectorum (Cynipis quercus-folii) punctione producta. Citò crescit Galla, et solitâ suâ magnitudine potitur, unius vel duorum dierum spatie, antequam insecta ovis excluduntur, quæ diurnâ irritatione tumoris incrementum efficiunt.

Larvis nondum in muscas mutatis, ne taberis interiora exeuntès devorent, decerpi debent Gallæ.

VIS. Gallæ amaræ, et astringentes, duræ, fragiles, latius striatæ: Tannin et Acido Gallico abundant.

USUS. In Gargarismatibus, injectionibus, et unguentis conficiendis.

180. QUERCUS PEDUNCULATA, (Racemosa.)

CHAR. SPEC. Foliis oblongis subsessilibus, glabris, sinuatis, lobis retundatis, fructibus oblongis pedunculatis.

ANGLICE, Common Oak.

LOCUS. In montibus, sylvis. Perennis.

VIS. Cortex amarus, astringens.

USUS. Decoctum pro injectione in Leucorrhœâ adhibitum.

CLASSIS XXI.

ORDO VIII.

MONGECIA. MONADELPHIA.

PINUS. Willd. Spec. Plant. iv. 494. Persoon Syn. ii. 578.

ORDO NAT. CONIFERÆ. Linn.

CLASSIS XV. ORDO V. CONIFERÆ. H. Calyx nullus. Squamæ staminiferæ. Coniferæ veræ. Jam.

CHAR. GEN. Masculus. Calyx tetraphyllus. Stamina plurima. Antheræ nudæ. Fœminicus. Calyces strobili sive conî: squama biflora. Pistillum unicum. Nux ala membranacea excepta.

* Foliis geminatis.

181. PINUS SYLVESTRIS.

CHAR. SPEC. Foliis geminis rigidis, conis ovato-conicis subgeminis junioribus pedunculatis recurvis.

ANGLICE, Scotch Fir.

LOCUS. Scotiæ montibus, et Europâ Septentrionali. Perennia.

Optimè se habet Pinus, annos septuaginta nata; post quadraginta autem Terebinthinam vulgarem præbet. Mense Maiâ ineunte, arbor ad sex unciarum spatium decorticata est, unde saccus resinusus pellucidus guttatim brevi exudat, et ad Septembrem usque ætia abunde defluere perstat. Hoc modo Pinus vegeta circiter decem Terebinthinæ libras quotannis suppeditare potest, per centum vices et ultra.

Viginti quinque libris destillatis, Olei Terebinthinæ circiter sex libras adipisci solent. Reliquum e Resinâ constat.

Ligni hujus arboris semicombustione, in frustula concisai, in annulis conservati, et cespiti tacti Pix liquida acquisita est.

182. PINUS LARIX.

CHAR. SPEC. Foliis fasciculatis deciduis strobilis ovato-oblongis squamarum marginibus reflexis laceris, bracteatis, pandaniformibus.

ANGLICE, Larch.

LOCUS. America, Europa Australior., Perennia.

Ex hac arbore, Terebinthinam Venetam, modo supradicto, obtinemus.

*** Foliis solitariis, et basi distinctes. Coni squamis levibus attenuatis.

183. PINUS BALSAMEA.

CHAR. SPEC. Foliis solitariis planis emarginatis sub-

pectinatis suprà suberectis, squamis coniflorentis acuminatis reflexis. Folia latiora, obtusiora, sæpè bifida.

LOCUS. In Virginie, Canadæ frigidissimis.

Eodem modo ex hac Pino Trebinthinâ Canadensi fruimus. Perennis.

184. PINUS ABIES.

CHAR. SPEC. Foliis solitariis tetragonis, strobilis cylindricis, squamis rhombeis complanatis margine repandis erosis.

ANGLICE, Norway Spruce Fir.

LOCUS. In Europæ, et Asiæ Borealibus sylvis montosis. Perennis.

Ex hac arbore prodit Pix Burgundica.

USUS. Terebinthinæ, et earum olea destillata, stimulantem, catharticam, diureticam anthelminticam vim habent; et extrinsecus ad ruborem excitandum applicantur. Intus præcipuè, Veneta et Canadensis exhibentur.

Largiore quantitate intestina, exigua renes maximè afficiuntur; quorum secretioni odorem violaceum impertiunt; et hæc vis tum subtili naturâ gaudet, ut alicujus urina cameram ineuntis, ubi, olei terebinthinæ vapor proderit, intra horæ quadrantem violam apertè redolere solet.

Merito laudatæ sunt Terebinthinæ in Gonorrhœâ, Leucorrhœâ, Lumbagine, Sciaticâ, Rheumatismo; et in Tœniis eliminandis.

DOSIS. Terebinthinarum gr. x. ad ʒi. Olei destillati ʒss. ad ʒj. Pro emplastris conficiendis nonnunquam usurpantur.

COMP. Linimentum Terebinthinæ. *Phar. Eblan.* Emplastrum Resinæ. *Phar. Lond. Edin. et Eblan.* Ceratum Resinæ. *Lond. et Edin.* Emp. Picis Comp. *Lond. et Edin.* Aqua Picis liquidæ. *Eblan.* Unguentum Picis liquidæ. *Eblan.*

CROTON. Willd. Spec. Plant. iv. 531. Persoon Syn. ii. 583.

ORDO NAT. TRICOCCÆ. Linn.

CLASSIS XV. ORDO I. EUPHORBLÆ. Sectio I. Juss.

CHAR. GEN. Masculus. Calyx cylindricus, quinque dentatus. Corolla pentapetala, (passim nulla.) Stamina decem ad quindecim, (passim libera.) Fœmineus. Calyx penta-sive polyphyllus. Corolla nulla. Styli tres, bifidi. Capsula tricocca, trilocularis trisperma.

185. CROTON TIGLIUM.

CHAR. SPEC. Foliis ovatis acuminatis, serratis, glabris basi bi-glandulosis; petiolis folio brevioribus, racemis terminalibus.

ANGLICE, Purging Croton.

LOCUS. In Moluccis Insulis. Perennis.

VIS. Stirpis singula pars drastica, sed semen præcipue valet.

DOSIS. Olei e seminibus expressi minimum unicum sacchari frustulo instillatum dosis ampla.

USUS. In Trismo, Apoplexiâ, Epilepsiâ, Maniâ et Encephali injuriis, ut Catharticum præstat: bene tamen cavendum, ne in Phlogisticis intestinorum morbis adhiberetur, et funesta, ne dicamus lethalia, orirentur.

186. CROTON ELUTERIA.

CHAR. SPEC. Foliis ovatis acuminatis integerrimis glabris subtus argenteo-squamulosis, racemis compositis axillaribus, caule arboreo.

ANGLICE, Cascarilla.

LOCUS. Bahamæ Insulæ, Jamaica. Perennis.

VIS. Fragrans, amara, aromatica.

USUS. Febribus, Dyspepsiâ, &c.

COMP. Inf. Cascarilla. *Phar. Lond.* Tinct. Cascarilla. *Phar. Lond.* et *Eblan.* Ext. Cascarilla. *Phar. Eblan.*

RICINUS. Willd. Spec. Plant. iv. 564. Persoon Syn. ii. 587.

ORDO NAT. TRICOCCÆ. Linn.

CLASSIS XV. ORDO I. EUPHORBIÆ. Sectio I.
Juss.

CHAR. GEN. Masculus. Calyx quinquepartitus. Stamina numerosa. Fœmineus. Calyx tripartitus. Styli tres, bifidi. Capsula ut plurimum echinata, trilocularis trisperma.

* Foliis palmatis.

187. RICINUS COMMUNIS.

CHAR. SPEC. Foliis peltatis palmatis, lobis lanceolatis serratis, caule herbaceo pruinoso.

ANGLICE, Castor Oil Plant.

LOCUS. Indiæ Occidentalis Insulæ, et America Australior. Annua.

VIS. Oleum e seminibus expressum vim benè laxativam exercet. Seminis corculum acerrimam ventriculi inflammationem, syncopen, et alia molestia, necnon lethalia, inducit. Absimile plurimis, oleis fixis, cum Alkohole facilè coalescit, unde integritas ejus commodè probatur.

DOSIS. ʒss. ad ʒj.

MOMORDICA. Willd. Spec. Plant. iv. 601. Persoon Syn. ii. 592.

ORDO NAT. CUCURBITACEÆ. Linn.

CLASSIS XV. ORDO II. CUCURBITACEÆ.

III. Stylus unicus. Fructus multilocularis polyspermus.
Juss.

CHAR. GEN. Masculus. Calyx quinquefidus. Corolla quinquepartita. Filamenta tria. Fœmineus. Stylus trifidus. Pepo sicca elasticè distiliens. Semina compressa.

188. MOMORDICA ELATERIUM.

CHAR. SPEC. Pomis ellipticis hispidis, foliis cordatis hispidis obtusis dentatis, cirrhis nullis.

ANGLICE, Wild Cucumber.

LOCUS. In Italiâ, Narbonâ, Siciiliâ Annua.

Peponium Elaterii Succus, qui tam acris naturæ est, ut cutem inflammare et exulcerare possit, post paucas horas

levem farinam cinereo-viridem demittit, quæ Elaterii Extractum vulgo appellatur.

USUS. In Anasarca, Ascitide, et Hydrothoraci; quando ægrotantis vires tam forte remedium sustinere valent.

DOSIS. Elaterii sinceri granum dimidium vel unicum, nauseam primò excitat, deinde vi Drasticâ et Hydrogogâ potitur.

CACUMIS. Willd. Spec. Plant. iv. 611. Persoon Syn. ii. 593.

ORDO NAT. CUCURBITACEÆ. Linn.

CLASSIS XV. ORDO II. CUCURBITACEÆ. Sectio III. Juss.

CHAR. GEN. Masculus. Calyx quinquedentatus. Corolla quinque partita. Filamenta tria. Fœmineus. Stigmata tria, crassa, bipartita. Peponis semina arguta, gemino ordine.

189. CUCUMIS COLOCYNTHIS.

CHAR. SPEC. Folijs multifidis, pomis globosis glabris.

LOCUS. Ad Bonæ Spei Promontorium. Annua.

VIS. Amarissima. Pulpa ejus drastica, dejectiones sanguineas interdum causat, et nunquam per se adhibetur.

COMP. Ext. Colocynthis. *Lond.* Ext. Colocynthis Comp. *Lond.* et *Eblan.* Pilulæ Aloës cum Colocynthide. *Eblan.*

CLASSIS XXII.

ORDO II.

DIECIÆ. DIANDRIA.

SALIX. Willd. Spec. Plant. iv. 653. Persoon Syn. ii. 598.

ORDO NAT. AMENTACEÆ. Linn.

CLASSIS XV. ORDO IV. AMENTACEÆ.

II. Flores dioici. Juss.

CHAR. GEN. Masculus. Amentum cylindraceum.
 Calyx squama. Glandula baseos nectarifera.
 Fœmineus. Flos maris. Stylus bifidus. Capsula unilobularis, bivalvis. Semina papposa.

* Foliis glabris serratis.

190. SALIX FRAGILIS.

CHAR. SPEC. Foliis lanceolatis acuminatis glabris glanduloso-serratis, amentis coætaneis, nectario masculino gemino, germinibus subsessilibus, lanceolatis glabris.

ANGLICE, Crack-willow.**LOCUS.** Angliæ indigena. Perennis.**VIS.** Amara.

*** Foliis villosis.

191. SALIX CAPREA.

CHAR. SPEC. Foliis ovatis acuminatis serratis undulatis subtus tomentosis, stipulis sublunatis, capsulis ventricosiss.

ANGLICE, Great Round-leaved Willow.**LOCUS.** Angliæ indigena. Perennis.**VIS.** Astringens, amara.**USUS.** In Dyspepsiâ olim.**DOSIS.** 3ss, ad ʒi.**192. SALIX ALBA.**

CHAR. SPEC. Foliis lanceolatis acuminatis serratis utrinque pubescentibus: serraturis infimis glandulosis.

ANGLICE, White Willow.**LOCUS.** Angliæ indigena. Perennis.**VIS et USUS.** Præcedentis.

CLASSIS XXII.

ORDO V.

DICECIA. PENTANDRIA.

HUMULUS. Willd. Spec. Plant. iv. 769. Persoon Syn. ii. 618.

ORDO NAT. SCABRIDÆ. Linn.

CLASSIS XV. ORDO III. URTICÆ.

Sectio II. Juss.

CHAR. GEN. Masculus. Calyx pentaphyllus. Corolla nulla. Fœmineus. Calyx monophyllus, obliquè patens, integer. Styli duo. Semen unicum intra calycem foliatum, (strobilum.)

193. HUMULUS LUPULUS.

ANGLICE, Hop.

LOCUS. Angliæ indigena. Perennis.

VIS. Amara, narcotica?

USUS. Rheumatismo, &c.

DOSIS. Ext. gr. iv. Tinct. 3i.

COMP. Ext. Humuli. *Lond.* Tinct. Humuli. *Lond. Edin.*

PISTACIA. Willd. Spec. Plant. iv. 750. Persoon Syn. ii. 615.

ORDO NAT. AMENTACEÆ. Linn.

CLASSIS XIV. ORDO XII. TEREINTACEÆ.

Sectio II. Juss.

CHAR. GEN. Masculus. Calyx quinquefidus. Corolla nulla. Fœmineus. Calyx trifidus. Corolla nulla. Styli quinque. Drupa monosperma.

194. PISTACIA TEREINTHUS.

CHAR. SPEC. Foliis impari-pinnatis, foliolis subseptenis ovato-lanceolatis basi rotundatis acutis mucronatis.

ANGLICE, Chio Turpentine.

LOCUS. Chio, et Cyprus Insulæ. Perennis.

VIS. Terebinthina Chia fragrans, stimulans.

195. PISTACIA LENTISCUS.

CHAR. SPEC. Foliis abruptè pinnatis, foliolis lanceolatis octonis, petiolo alato.

ANGLICE, Mastic.

LOCUS. Chios Insula. Perennis.

VIS. Fragrans.

CLASSIS XXII.

ORDO VI.

DIOECIA. HEXANDRIA.

SMILAX. Willd. Spec. Plant. iv. 773. Persoon Syn. ii. 618.

ORDO NAT. SARMENTACEÆ Linn.

CLASSIS III. PLANTÆ MONOCOTYLEDONES.

STAMINA PERIGYNA. ORDO II. ASPARAGI.

II. Flores dioici. Germen superum. Juss.

CHAR GEN. Masculus. Calyx pentaphyllus. Corolla nulla. Fœmineus ut in mare. Styli tres. Bacca triloculatis. Semina duo.

* Caule aculeato angulato.

196. SMILAX SARSAPARILLA.

CHAR. SPEC. Caule subtetrageno, foliis intermixtis ovato lanceolatis, cuspidatis subquinque nerviis, subtus glaucescentibus.

ANGLICE, Sarsaparilla.

In LOCUS. In America Meridionali, Virginia, India Occidentali. Perennis.

VIS. Cortex amaricans. Pars interior, farinâ mucilaginosa abundat. Varietas rubra e Jamaicâ deportata, præstantior existimata.

USUS. Temporibus variis Sarsaparilla in usum recepta, et iterum prætermissa fuit. Nuperius frequentius adhibita est, et largior extracti liquidi dosis in cutis eruptionibus, Cynanche, et aliis morbis Syphiliticis feliciter usurpari videtur.

DOSIS. Pulv. ʒi. ad ʒiij. Decoct. ʒiv. ad ʒviii.

COMP. Decoctum Sarsaparillæ. *Lond. Edin. et Eblan.*
Dec. Sarsaparillæ Comp. *Lond. Eblan.* Extractum Sarsaparillæ. *Lond.*

CLASSIS XXII.

ORDO X.

DICECIA. DODECANDRIA.

MENISPERMUM. Willd. Spec. Plant. iv. 824. Persoon Syn. ii. 627.

ORDO NAT. SARMENTACEÆ. Linn.

CLASSIS XIII. ORDO XVII. MENISPERMA.
Juss.

CHAR. GEN. Masculus. Calyx diphyllus. Petala quatuor sive sex exteriora, octo interiora. Stamina sedecim. Fœmineus. Corolla maris. Stamina octo sterilia. Germina duo sive tria. Baccæ binæ, monospermæ. (Cotyledones segregatæ.)

197. MENISPERMUM PALMATUM.

CHAR. SPEC. Piloso-hispidum, foliis quinquelobis, plerisque palmatis, basi cordatis, lobis acumınatis.

ANGLICE, Calumba.

LOCUS. Africa Australior. Perennis.

VIS. Amara, aromatica.

USUS. In Dyspepsiâ, &c.

DOSIS. Pulv. gr. xv. ad 3ss. ter quaterve die.

COMP. Infusum Calumbæ. *Lond. Edin.* Tinctura Calumbæ. *Lond. Edin.* et *Eblan.*

CLASSIS XXII.

ORDO XIII.

DICECIA. MONADELPHIA.

JUNIPERUS. Willd. Spec. Plant. iv. 851.

Persoon Syn. ii. 632.

ORDO NAT. CONIFERÆ. Linn.

CLASSIS XV. ORDO V. CONIFERÆ. Sectio II.

Juss.

CHAR. GEN. Masculus. Amentum ovatum. Calyx squama, Stamina tria. Fœmineus. Calyx tripartitus. Petala tria. Styli tres. Bacca trisperma, tribus tuberculis calycis inæqualis. (Nux ossea, unilocularis basi glandulosa.)

198. JUNIPERUS SABINA.

CHAR. SPEC. Foliis oppositis obtusis medio glandulosis, quadrifariam imbricatis, tenellis acutis oppositis, caule fruticoso.

ANGLICE, Savin.

LOCUS. In Lusitaniâ, Italiâ, Siberiâ. Perennis.

VIS. Acris graveolens, amara, oleum essenziale continens, stimulans, emmenogoga, abortum inducens, anthelmintica?

USUS. Amenorrhœa; extrinsecus Verrucis, applicatur, et, ad vesicatorii inflammationem excitandam, ne vulnere induceretur cicatrix.

DOSIS. Gr. v. ad x.

COMP. Oleum Volatile Juniperi Sabinæ. *Edin. et Eblan.*
Ext. Sabinæ. *Eblan.* Ceratum Sabinæ. *Lond.*

199. JUNIPERUS COMMUNIS.

CHAR. SPEC. Foliis ternis patentibus mucronatis, baccâ longioribus.

ANGLICE, Juniper.

LOCUS. Angliæ indigena. Perennis.

VIS. Dalcis, aromatica, pungens, diuretica.

USUS. In Hydrops.

COMP. Oleum Juniperi. *Lond. Edin. et Eblan.* Spiritus Janip. Comp. *Lond. Edin. et Eblan.*

MYRISTICA. Willd. Spec. Plant. IV. 869. Persoon Syn. ii. 635.

CLASSIS VI. ORDO IV. LAURI. Genera Lauris affinia. Juss.

CHAR. GEN. Masculus. Calyx nullus. Corolla campanulata, trifida. Filamentum columnare. Antheræ sex ad decem, connatæ. Fœmineus ut in mare. Stylus nullus. Stigmata duo. Drupa nuce arillata, monosperma. Semen magnum, intus venoso-variegatum.

200. MYRISTICA MOSCHATA.

CHAR. SPEC. Foliis oblongis acuminatis glabris, venis simplicibus, fructu solitario glabro.

ANGLICE, Nutmeg.

LOCUS. In Moluccis Insulis. Perennis.

Tegumentum seminis externum, Anglicè, *Mace.*

VIS. Fragrans, aromatica, oleum essentielle continens. Purgantia, et alia corrigens.

DOSIS. Gr. v. ad x.

COMP. Spiritus Myristicæ. *Lond. Edin. et Eblan.* Conf. Aromatica.

CLASSIS XXIII.

ORDO I.

POLYGAMIA. MONŒCIA.

VERATRUM. Willd. Spec. Plant. iv. 895. Persoon Syn. i. 397.

CLASSIS III. ORDO III. JUNC. IV. Germina plura (sæpius tria.) Capsulæ totidem uniloculares, interdum basi coalitæ. Flores paniculati aut spicati. Juss.

CHAR. GEN. Hermaphroditus. Calyx nullus. Corolla hexapetala. Stamina sex. Pistilla tria. Capsulæ tres polyspermæ. Masculus. Calyx nullus. Corolla hexapetala. Stamina sex. Pistilli rudimentum.

201. VERATRUM ALBUM.

CHAR. SPEC. Racemis paniculatis, bracteis ramorum oblongis partialibus pedunculum pubescentem subæquantibus floribus erectis.

ANGLICE, White Hellebore.

LOCUS. Siberiæ, Austriæ, Helvetiæ, Græciæ, montosis. Perennis.

VIS. Alkali continet hæc radix, quod DD. Pelletier et Caventou nuper in lucem sub nomine Veratriæ ediderunt, et cui Colchici radicis tribuuntur vires.

Memoratu dignum est has duas plantas Veratrum, et Colchicum, in Systemate Naturali D. Jussieu, juxta positas inveniri.

Temporibus priscis in Maniâ, Epilepsiâ et Paralysi adhiberi solebat Veratrum; sed in hodiernâ praxi rarò usurpatur ob actionem ejus acrem et ancipitem. Dosis enim modica vomitum, catharsin, syncopen, et taliâ periculosa infert. Decoctum extrinsecùs applicatum Scabiæ lymphatica, et aliis, naturâ simili, morbis bene convenit. Minimè tamen vel extrinsecùs, incautè et temerè, tractandum.

STALAGMITIS. Willd. Spec. Plant. iv. 980. Persoon Syn. ii. 68.

ORDO NAT. TRICCOCCÆ. Linn.

CLASSIS XIII. ORDO IX. GUTTIFERÆ. I. Stylus nullus. Juss.

CHAR. GEN. Hermaphroditus. Calyx tetraphyllus. Corolla tetrapetala. Stamina triginta. Receptaculo quadrangulo carnosio inserta. Stylus crassus. Stigma quadrilobum. Bacca unilocularis stylo coronata trisperma.

Masculi. Calyx, corolla, et stamina hermaphrodita.

202. STALAGMITIS CAMBOGIOIDES.

ANGLICE, Camboge.

LOCUS. Zeylona. Perennis.

VIS. Vi drastica potitur hoc gummi, morbis plurimis convenienti. Granorum paucorum dosis, ad Tænias vermes expellendos maximè valet, sed vomitus, et tormina hand rarò consequuntur.

ACACIA. Willd. Spec. Plant. iv. 1049.

ORDO NAT. LOMENTACEÆ. Linn.

CLASSIS XIV. ORDO XI. LEGUMINOSÆ.

Sectio I. Juss.

CHAR. GEN. Hermaphroditus. Calyx quinquedentatus. Corolla quinquefida vel pentapetala. Stamina quatuor ad centum. Pistillum unicum. Legumen bivalve.

Masculi. Calyx quinquedentatus. Corolla quinquefida seu pentapetala. Stamina quatuor ad centum.

**** Foliis duplicato pinnatis, spinis sive aculeis stipularibus spicis elongatis.

203. ACACIA CATECHU.

CHAR. SPEC. Aculeis geminis stipularibus uncinatis, foliis bipinnatis, partialibus decemjugis, propriis multijugis pubescentibus, glandula petiolari et inter duo terminali-partialium, spicis cylindræis geminis ternisve axillaribus.

ANGLICE, Catechu.

LOCUS. Malabaria, Zeylona. Perennis.

VIS. Principiis abundat Catechu astringentibus, Tannin et Acido Gallico, nuncupatis; ideoque Diarrhœâ, et aliis

morbis affinis datur. Gargarismatis formâ interdum ad Uvulam laxatam reficiendam adhibetur. Cantores vocis raucitate laborantes, extracti catechu modicum manducare dicuntur.

***** Foliis duplicato pinnatis spinis sive aculeis stipularibus spicis globosis.

204. ACACIA VERA.

CHAR. SPEC. Spinis stipularibus geminis patentibus, foliis bipinnatis, partialibus bijugis, propriis octo-decjugisve, glandula inter singula partialium paria, spicis subgeminis pedunculatis axillaribus.

ANGLICE, Gum Arabic.

LOCUS. Arabia Petrea, Ægyptus. Perennis.

VIS et USUS. Gummi Demulcens dictum. Nutriens.

Anno 1750, secundum D. Hasselquist, Mercatores ex Abyssiniâ congregati iter conficientes, cibo omni consumpto, et morte mox iis impendente, Acaciæ gummi copiam inter merces fortè reperiere, quo solo hominum mille per duas menses vitam sustinere.

CLASSIS XXIII.

ORDO II.

POLYGAMIA. DICECIA.

FRAXINUS. Willd. Spec. Plant. iv. 1098.

ORDO NAT. SEPIARIÆ. Linn.

CLASSIS VIII. ORDO IV. JASMINEÆ. I. Fructus capsularis. Juss.

CHAR. GEN. Hermaphroditi. Calyx nullus sive quadripartitus. Corolla nulla sive tetrapetala. Stamina duo Pistillum unicum. Samara monosperma ala lanceolata terminata.

Fœminei. Calyx nullus sive quadripartitus. Corolla

nulla sive tetrapetala. Pistillum unicum. Samara monosperma ala lanceolata terminata.

ORNUS EUROPÆA. Persoon Syn. i. 9.

205. FRAXINUS ORNUS.

CHAR. SPEC. Foliolis petiolatis oblongo-lanceolatis acuminatis serratis, floribus corollatis.

ANGLICE, Flowering Ash.

LOCUS. Calabriâ, Siciliâ. Perennis.

VIS. Manna, quæ ex hac plantâ prodit, laxativa.

DOSIS. ʒj. ad ʒi.

FICUS. Willd. Spec. Plant. iv. 1131. Persoon Syn. ii. 608.

ORDO NAT. SCABRIDÆ. Linn.

CLASSIS XV. ORDO III. URTICÆ. Sectio I. Juss.

CHAR. GEN. Receptaculum commune carnosum, squamis connivens, occultans flosculos apetalos, vel in eodem vel distincto individuo.

Masculi. Calyx tripartitus.

Fœminei. Calyx quinquepartitus. Pistillum unicum, laterale. Semen unicum calyce persistenti clauso subcarnoso tectum.

* Foliis lobatis.

206. FICUS CARICA.

CHAR. SPEC. Foliis cordatis tri-quinquelobisve repando-dentatis, lobis obtusis, suprâ scabris, subtus pubescentibus.

ANGLICE, Fig Tree.

LOCUS. Græciâ, Hispaniâ, Italiâ. Perennis.

VIS. Laxativa.

CLASSIS XXIV.

ORDO I.

CRYPTOGAMIA. FILICES.

Herba frondosa. Fructificatio vel dorsalis, vel terminalis, vel subradicalis.

* Annulatæ; capsulis annulo elastico, valvulis contrario, cinctis.

ASPIDIUM. Willd. Spec. Plant. v. 211. Smith Fl. Brit. iii. 1118.

ORDO NAT. FILICES. Linn.

CLASSIS I. PLANTÆ ACOTYLEDONES. ORDO V. FILICES. II. Folliculi frondibus subtus impositi. Cætera ignota. Juss.

CHAR. GEN. Fructificationes in punctis, subrotundis sparsis non marginalibus. Involucrum umbilicatum, undique ferè debiscens.

** Fronde subbipinnatâ.

207. ASPIDIUM FILIX MAS.

CHAR. SPEC. Fronde bi-pinnata: pinnulis obtusis serratis stipite paleaceo involucris circinatis.

ANGLICE, Male Fern.

LOCUS. Angliæ sylvis. Perennis.

VIS. Anthelmintica existimata.

CLASSIS XXIV.

ORDO IV.

CRYPTOGAMIA. LICHENES.

LICHEN. Linn. Syst. Vegetab. curâ Persoon. 995.

ORDO NAT. ALGÆ. Linn.

CLASSIS I. ORDO II. ALGÆ III. Coriaccæ aut crustacæ. Fructificatio spectabilior. Lichenes. Juss.

CHAR. GEN. Flores Masculi. Receptaculum subrotundum, planiusculum, nitidum. Flores Fœminei. Farina foliis aspersa.

D. * Foliacei.

208. LICHEN ISLANDICUS.

CETRARIA ISLANDICA. Acharii Methodus. 293.

CHAR. SPEC. Foliaceus ascendens laciniatus: marginibus elevatis ciliatis. Linn. Spec. Plant. 1611. English Botany. 1330.

ANGLICE, Iceland Moss.

LOCUS. Angliæ indigena.

VIS. Amara, mucilaginosa.

USUS. Olim in Phthisi, Diarrhœâ, Dysenteria.

COMP. Dec. Lichenis. *Lond. et Eblan.*

***** Fruticulosi.

209. LICHEN ROCCELLA.

CHAR. SPEC. Lichen fruticulosus solidus aphyllus subramosus tuberculis alternis. Dickson. Plant Cryptogam. Britannicæ. Fasc. iii. 19.

ANGLICE, Orchall, or Dyers' Lichen.

LOCUS. In Insulæ Guernsey rupibus marinis.

VIS. Violacea-tinctoria. Tinctoriis et Philosophis Chemicis usurpatur.

CLASSIS XXIV.

ORDO V.

CRYPTOGAMIA. ALGÆ.

FUCUS. Linn. Syst. Vegetab. curâ D. Persoon. p. 1004. Linn. Genera Plant. curâ D. Schreber, tom. ii. p. 768.

ORDO NAT. ALGÆ. Linn.

CLASSIS I. ORDO II. ALGÆ. II. Membranaceæ
aut coriaceæ. Fructificatio incerta. Fuci. Juss.

CHAR. GEN. * Masculi flores? Vesiculæ glabræ,
cavæ, pilis intus adpersæ.

* Fœminei flores.

Vesiculæ glabræ, gelatinâ repletæ, adpersæ punctis
perforatis semine foetis. Semina solitaria.

** Alati sive Fronde planâ, stipite medium folium percar-
rente.

210. FUCUS VESICULOSUS.

CHAR. SPEC. Fronde dichotomâ integerrimâ; vesi-
culis innatis axillaribusque; apicibus tumidis tuberculatis
acutiusculis. DD. Goodenough et Woodward in Societa-
tis Linn. Actis tom. iii. p. 103.

ANGLICE, Bladder Fucus.

LOCUS. In litoribus.

VIS. Stimulans, mucilaginosa.

USUS. Tumoribus glandulosis planta recens quasi cata-
plasma stimulans olim usurpata est.

Fuci Vesiculosi cineres, quantitatem Iodinæ exiguam for-
sân continentes, pro Spongiâ ustâ contra Scrophulam et
Bronchocelen medicinæ praxi adhiberi solebant.

Regionibus quibusdam ad porcos etiam alendos, et agros
fœcundandos, planta recens sæpè inservit.

Combustione ejus, in cumulos litoribus coacervatæ, magna
cinerum copia obtinetur. Hæc materia, *Kelp* Anglicè
dicta, Sodæ salibus scatet, idcirco ad saponem durum et
vitrum conficiendum largè usurpatur.

CLASSIS XXIV.

ORDO VI.

CRYPTOGAMIA. FUNGI.

BOLETUS. Linn. Gen. Plant. curâ Schreber, tom. ii. 769.

ORDO NAT. FUNGI. Linn.

CLASSIS I. ORDO I. FUNGI. II. Fungi carnosii pileati, pileis ~~orbicularibus, sessilibus aut saepe~~ stipiti centrali impositis.

CHAR. GEN. Fungus horizontalis, subtile porosus.

* Parasitici acaules.

211. BOLETUS IGNIARIUS.

CHAR. SPEC. Acaulis pulvinatus laevis, poris tenuissimis. Linn. Spec. Plant. ii. 1645.

LOCUS. Indigena, arborum truncis.

USUS. Olim ad Hæmorrhagiam cohibendam.

SYNOPSIS
PLANTARUM MEDICINALIUM,
SECUNDUM SYSTEMA,
D. JUSSIEU, DISPOSITARUM.

CLASSIS I.

PLANTÆ ACOTYLEDONES.

ORDO I. FUNGI. Sectio II. Fungi carnosī pileatī, pileis orbicularibus, sessilibus aut sæpius stipiti centrali impositis.

Boletus.

ORDO II. ALGÆ. Sectio II. Membranaceæ, aut coriaceæ. Fructificatio incerta.

Fucus.

Sectio III. Coriaceæ, aut crustaceæ. Fructificatio spectabilior.

Lichen.

ORDO V. FILICES. Sectio II. Folliculi frondibus subtus impositi. Cætera ignota.

Aspidium.

CLASSIS II.

PLANTÆ MONOCOTYLEDONES. STAMINA
HYPOGYNA.

ORDO I. AROIDEÆ.

Sectio I. Spadix spathâ involutus.

Arum.

Sectio II. Spadix nudus, spathâ destitutus.

Acorus.

ORDO IV. GRAMINEÆ. Sectio II. Styli duo. Stamina tria. Gluma uniflora.

Saccharum.

Sectio VII. Styli duo. Stamina tria. Glumæ multifloræ, supra axim seu rachim densè spicatæ. Spica sæpiùs simplex.

Hordeum.

Triticum.

Secale.

Sectio VIII. Styli duo. Stamina tria. Glumæ multifloræ vagæ. Flores sæpiùs paniculati.

Avena.

CLASSIS III.

PLANTÆ MONOCOTYLEDONES. STAMINA PERIGYNA.

ORDO I. PALMÆ. Sectio I. Frondes pinnatæ. Foliola sæpiùs lanceolata angusta.

Cocos.

ORDO II. ASPARAGI. Sectio II. Flores dioici. Germen superum.

Smilax.

ORDO III. JUNCII. Sectio IV. Germina plura (sæpiùs tria.) Capsulæ totidem uniloculares, interdum basi coalitæ. Flores paniculati aut spicati.

Veratrum.

Colchicum.

ORDO VI. ASPHODELI. Sectio I. Flores spicati. Radix fibrosa. Calyx tubulosus.

Aloe.

Sectio IV. Flores spicati. Radix bulbosa. Calyx sex-partitus, basi staminifer.

Scilla.

Sectio V. Flores umbellati. Radix bulbosa. Calyx sex-partitus æqualis.

Allium.

ORDO VIII. IRIDES. Sectio II. Stamina filamentis distinctis.

Iris.

Crocus.

ORDO IX. BORAGINEÆ.

Sectio IV. Fructus gymno-tetraspermus. Faux corollæ instructa quinque squamis calcarum instar cavis, intra corollam prominulis in ejusdem laciniarum basi, extrâ hiulcis. Herbæ plerumque asperifoliæ.

Anchusa.

ORDO X. CONVULVULI. Sectio I. Stylus unicus.
Convolvulus.

ORDO XIII. GENTIANÆ.

Sectio I. Capsula simplex unilocularis.

Chironia. (*Erythræa.*)

Gentiana.

Sectio II. Capsula simplex bilocularis.

Sectio III. Capsula didyma bilocularis.

Spigelia.

ORDO XIV. APOCINEÆ.

Sectio IV. Genera Apocineis affinia, non lactescentia.

Strychnos.

CLASSIS IX.**PLANTÆ DICOTYLEDONES MONOPETALÆ.
COROLLA PERIGYNA.**

ORDO I. GUAIACANÆ. Sectio I. Stamina definita.

Styrax.

ORDO II. RHODODENDRA.

Sectio I. Corolla Monopetala.

Rhododendron.

ORDO III. ERICÆ. Sectio I. Germen superum.

Arbutus.

CLASSIS X.**PLANTÆ DICOTYLEDONES MONOPETALÆ.
COROLLA EPIGYNA. ANTHERÆ CONNATÆ.**

ORDO I. CICHORACEÆ. Sectio II. Receptaculum nudum. Semen papposum, pappo piloso.

Lactuca.

Leontodon.

ORDO II. CINAROCEPHALÆ. Sectio I. Cinarocephalæ veræ. Squamæ calycis spinosæ.

Arctium.

Centaurea.

ORDO III. CORYMBIFERÆ. Sectio II. Receptaculum nudum. Semen papposum. Flores radiati.

Solidago.

Inula.

Tussilago.

Arnica.

Sectio IV. Receptaculum nudum. Semen nudum seu non papposum. Flores flosculosi.

Tanacetum.

Artemisia.

Sectio V. Receptaculum paleaceum. Semen nudum seu non papposum. Flores plerumque radiati, rarò flosculosi.

Anthemis.

CLASSIS XI.

DICOTYLEDONES MONOPETALÆ. COROLLA EPIGYNA. ANTHERÆ DISTINCTÆ.

ORDO I. DIPSACEÆ. Sectio II. Flores distincti.

Valeriana.

ORDO II. RUBIACEÆ. Sectio I. Fructus dicoccus dispermus. Stamina sæpiùs quatuor. Folia plerumque verticillata, caulis plerumque herbaceus.

Rubia.

Sectio IV. Fructus monocarpus bilocularis polyspermus. Stamina quinque. Folia opposita, caulis sæpè frutescens.

Cinchona,

ORDO III. CAPRIFOLIA. Sectio III. Calyx bracteatus. Stylus nullus, stigmata tria. Corolla monopetala.

Sambucus.

CLASSIS XII.

PLANTÆ DICOTYLEDONES POLYPETALÆ. STAMINA EPIGYNA.

ORDO II. UMBELLIFERÆ. Sectio I. Umbelliferæ veræ. Umbellæ et umbellulæ sæpiùs nudæ.

Pimpinella.

Carum.

Anethum.

• Pastinaca.

Sectio II. Umbelliferæ veræ. Umbellæ nudæ; umbellulæ involucrellatæ.

Coriandrum.

Sectio III. Umbelliferæ veræ. Umbellæ et Umbellulæ involucrellatæ.

Cuminum.

• Bubon.

Angelica.

Sium.

Ferula.

Conium.

• Daucus.

Sectio IV. Umbelliferæ anomalæ.

Eryngium.

CLASSIS XIII.

PLANTÆ DICOTYLEDONES POLYPETALÆ.

STAMINA HYPOGYNA.

• ORDO I. RANUCULACEÆ. Sectio II. Capsulæ polyspermæ intus dehiscentes. Petala irregularia. Calyx sæpè coloratus, a Linnæo corolla dictus, petalis ab eodem in nectaria conversis.

Helleborus.

Delphinium.

Aconitum.

• ORDO II. PAPAVERACEÆ. Sectio I. Stamina indefinita; antheræ filamentis adnatæ.

Papaver.

• ORDO III. CRUCIFERÆ. Sectio I. Fructus siliquosus. Stylus nullus.

Sinapis.

Cardamine.

Sectio II. Fructus siliculosus. Stylus unicus.

Cochlearia.

• ORDO VI. ACERA. Sectio I. Fructus multilocularis.

Æsculus.

• ORDO IX. GUTTIFERÆ. Sectio I. Stylus nullus.

Cambogia. (Stalagmitis.)

ORDO X. AURANTIA. Sectio II. Fructus poly-
permus baccatus. Folia pellucido-punctata. Aurantia vera.
Citrus.

ORDO XI. MELIÆ. Sectio I. Folia simplicia.

Wintera. (Canella.)

Sectio III. Genera Meliis affiniâ.

Swietenia.

ORDO XII. VITES.

Vitis.

ORDO XIII. GERANIA.

Genera Geraniis Affinia.

Oxalis.

ORDO XIV. MALVACEÆ. Sectio II. Stamina in
tubum corolliferum connata, indefinita. Fructus multicapsu-
laris; capsulæ verticillatæ, in orbem dispositæ aut in unam
compactæ.

Malva.

Althæa.

ORDO XV. MAGNOLIÆ.

Genera Magnoliis Affinia.

Quassia.

ORDO XVII. MENISPERMA.

Menispermum.

ORDO XX. CISTI.

Genera Cistis affinia fructu trivalvi, valvis seminiferis, sed
definitè staminifera.

Viola.

ORDO XXI. RUTACEÆ.

Sectio I. Folia stipulacea sæpiùs opposita.

Guaiacum.

Sectio II. Folia alterna nuda.

Ruta.

Sectio III. Genera Rutaceis affinia.

Diosma.

ORDO XXII. CARYOPHYLLÆ. Sectio V. Calyx
tubulosus. Stamina decem (alterna hypogyna, alterna
sæpiùs epipetala.) Styli duo, aut tres, aut quinque.

Dianthus.

Sectio VII. Genera Caryophylleis affinia.

Linum.

CLASSIS XIV.

PLANTÆ DICOTYLEDONES POLYPETALÆ. STAMINA PERIGYNA.

ORDO VII. MYRTI. Sectio I. Flores in foliorum axillis, aut in pedunculis multifloris oppositi. Folia plerumque opposita et punctata.

Melaleuca.

Myrtus.

Caryophyllus. (Eugenia.)

Punica.

ORDO IX. SALICARIÆ. Sectio I. Flores polypetali.
Lythrum.

ORDO X. ROSACEÆ. Sectio I. Germen simplex inferum, polystylum. Pomum calycino limbo umbilicatum, multiloculare. Arbores aut frutices. Pomaceæ.

Pyrus. (Cydonia.)

Sectio II. Germina plura indefinita, calyce urceolari supra coarctato tecta, quasi infero, singula monostyla. Semina totidem. Frutices.

Rosa.

Sectio III. Germina plura definita (rarius unicum,) calyce urceolari supra coarctato tecta, quasi infera, singula monostyla. Semina totidem. Herbæ pleræque, plures apetalæ, plures definitè staminiferæ, quædam diclines.

Agrimonia.

Sectio IV. Germina plura indefinita, verè supera, receptaculo communi imposita, singula monostyla. Semina totidem nuda aut rarius baccata. Herbæ aut rarius frutices.

Tormentilla.

Geum.

Sectio VII. Germen unicum superum monostylum.

Nux mono aut disperma, nuda aut sæpius drupacea.
Arbores aut frutices.

Prunus.

Amygdalus.

ORDO XI. LEGUMINOSÆ. Sectio I. Corolla regularis. Legumen multiloculare, sæpius bivalve, dissepimentis transversis, loculis monospermis. Stamina distincta. Arbores aut frutices, folia abruptè pinnata.

Mimosa. (Acacia.)

Tamarindus.

Cassia.

Sectio II. Corolla regularis. Legumen uniloculare bivalve. Stamina decem distincta. Arbores aut frutices. Folia abruptè pinnata.

Hæmatoxylon.

Sectio V. Corolla irregularis papilionacea. Stamina decem diadelpa. Legumen uniloculare bivalve. Frutices aut Herbæ: folia simplicia aut ternata aut rariùs digitata; stipulæ nunc subnullæ, nunc conspicuæ imo petiolo adnatæ, aut ab eodem distinctæ.

Spartium.

Dolichos. (Stizolobium.)

Sectio VI. Corolla irregularis papilionacea. Stamina decem diadelpa. Legumen uniloculare, bivalve. Herbæ aut frutices aut arbusculæ; folia impari-pinnata.

Astragalus.

Glycyrrhiza. (Liquiritia.)

Sectio IX. Corolla irregularis papilionacea. Stamina plerumque decemdiadelpa. Legumen capsulare uniloculare submonospermum sæpe non dehiscens. Arbores aut frutices; folia sæpius impari-pinnata; stipulæ a petiolo distinctæ mox deciduæ.

Geoffræa.

Pterocarpus.

Sectio X. Corolla irregularis interdum nulla. Stamina decem distincta. Legumen capsulare uniloculare submonospermum, sæpius non dehiscens. Arbores aut frutices; folia impari-pinnata aut simplicia; stipulæ a petiolo distinctæ mox deciduæ.

Copaifera.

ORDO XII. TEREINTACEÆ.

Sectio I. Germen simplex. Fructus unilocularis monospermus.

Rhus.

Sectio II. Germen simplex. Fructus multilocularis, loculis quibusdam interdum abortivis.

Amyris.

Pistacea.

ORDO XIII. Rhamni. Sectio III. Stamina petalis opposita. Fructus drupaceus.

Rhamnus.

CLASSIS XV.

PLANTÆ DICOTYLEDONES APETALÆ. STAMINA IDIOGYNA, SEU A PISTILLO SEGREGATA.

ORDO I. EUPHORBIÆ.

Sectio I. Styli plures definiti, sæpiùs tres.

Euphorbia.

Ricinus.

Croton.

ORDO II. CUCURBITACEÆ.

Sectio III. Stylus unicus. Fructus multilocularis polyspermus.

Momordica.

Cucumis.

ORDO III. URTICÆ.

Sectio I. Flores in communi involucro monophyllo reconditi.

Ficus.

Dorstenia.

Sectio II. Flores receptaculo communi multifloro impositi, aut squamis involucrantibus capitati, aut distincti sparsi.

Morus.

Humulus.

Sectio III. Genera Urticis affinia.

Piper.

ORDO IV. AMENTACEÆ.**Sectio I. Flores hermaphroditi.****Ulmus.****Sectio II. Flores dioici.****Salix.****Sectio III. Flores monoici.****Quercus.****ORDO V. CONIFERÆ.****Sectio II. Calyx nullus. Squamæ staminiferæ.****Coniferæ veræ.****Juniperus.****Pinus.****PLANTÆ INCERTÆ SEDIS.****MONOPETALÆ GERMINE SUPERO.****Galipea.****POLYPETALÆ GERMINE SUPERO.****Krameria.****APETALÆ DICLINES GERMINE SUPERO.****Myroxylon.**

INDEX GENERUM.

	Pag.			Pag.	
Acacia	119	135	Canella	64	133
Aconitum	74	132	Capsicum	25	129
Acorus	45	126	Cardamine	83	132
Æsculus	48	132	Carum	38	131
Agrimonia.	65	134	Caryophyllus	65	134
Allium	42	127	Cassia	55	135
Aloe	43	127	Cathartocarpus	56	
Alpinia	6	128	Centaurea	101	131
Althæa	86	133	Cetraria	122	126
Amomum	5	128	Chironia	22	130
Amygdalus	67	135	Cinchona	27	131
Amyris	48	136	Citrus	92	133
Anchusa	19	132	Cochlearia	83	132
Anethum	39	132	Cocos	104	127
Angelica	37	132	Colchicum	46	127
Anthemis	100	131	Conium	34	132
Arbutus	59	130	Convolvulus	21	130
Arctium	95	130	Copaifera	61	835
Aristolochia	102	128	Coriandrum	37	132
Arnica	98	131	Crocus	13	127
Artemisia	96	131	Croton	108	136
Arum	105	126	Cucumis	111	136
Asarum	63	128	Cuminum	35	132
Aspidium	121	126	Curcuma	5	128
Astragalus	90	135	Cydonia	68	134
Atropa	26	129			
Avena	14	127	Daphne	49	128
			Datura	22	129
Boletus	125	126	Daucus	34	132
Bubon	36	132	Delphinium	73	132
Butea	89		Dianthus	62	134
			Digitalis	82	129
Callicocca	29		Diosma	31	133
Cambogia	118	133	Dolichos	90	135

	Pag.		Pag.
Dorstenia	19 136	Melaleuca	93 134
Elettaria	6 128	Melissa	80 129
Eucalyptus	89	Menispermum	115 133
Eryngium	33 132	Mimosa	135
Erythraea	22 130	Mentha	77 129
Eugenia	65 134	Menyanthes	20 128
Euphorbia	102 136	Momordica	110 136
		Morus	103 136
Ferula	36 132	Myristica	117 128
Ficus	121 136	Myroxylon	54 137
Fraxinus	120 129	Myrtus	66 134
Fucus	123 126		
		Nicotiana	24 129
Galipea	8 137		
Gentiana	32 130	Olea	7 129
Geoffraea	92 135	Origanum	80 129
Geum	70 134	Ornus	120 129
Glycyrrhiza	91 135	Oxalis	62 133
Gratiola	9 129		
Guaiaacum	57 133	Papaver	71 132
		Pastinaca	38 132
Hæmatoxylon	56 135	Pimpinella	39 131
Helleborus	75 132	Pinus	106 137
Hordeum	17 127	Piper	10 136
Humulus	113 136	Pistacia	113 136
Hyoscyamus	23 129	Polygala	87 129
Hyssopus	76 129	Polygonum	50 128
		Prunus	68 135
Inula	98 131	Pterocarpus	88 135
Iris	13 127	Punica	66 134
		Pyrus	68 134
Juniperus	116 137		
Krameria	18 137	Quassia	58 133
		Quercus	105 137
Lactuca	94 130		
Laurus	51 128	Rhamnus	30 136
Lavandula	78 129	Rheum	53 128
Leontodon	94 130	Rhus	40 136
Lichen	122 126	Rhododendron	59 130
Linum	41 134	Ricinus	109 136
Liquiritia	91 135	Rosa	69 134
Lythrum	64 134	Rosmarinus	9 129
		Rubia	18 131
Malva	86 133	Rumex	47 128
Maranta	7 128	Ruta	58 133
Marrubium	79 129		
		Saccharum	14 126

	<i>Pag.</i>		<i>Pag.</i>
Salix	111 137	Tanacetum	97 131
Salvia	10 129	Tamarindus	85 135
Sambucus	40 131	Tormentilla	70 134
Scrophularia	81 129	Triticum	16 127
Scilla	15 127	Teucrium	79 129
Secale	44 127	Tussilago	99 131
Sinapis	86 132		
Sium	35 132	Ulmus	33 137
Smilax	114 127		
Solanum	26 129	Valeriana	12 131
Solidago	99 131	Viola	32 133
Spartium	89 135	Veratrum	118 127
Spigelia	20 130	Veronica	8 129
Stalagmitis	118 133	Vitis	31 133
Stizolobium	90 135		
Strychnos	24 130	Wintera	75 133
Styrax	60 130		
Swietenia	57 133	Zingiber	6 128


LONDINI:


EX TYPOGRAPHIA GULIELMI TYLSE,
BRIDGEWATER SQUARE.





3 2044 102 800 067







3 2044 102 800 067



